



SEBRING™ 2.65 ST

SEED TREATMENT FUNGICIDE

For use as a seed treatment for control of systemic downy mildew, Pythium seed rot, Pythium damping-off, and early season Phytophthora diseases of certain crops.

ACTIVE INGREDIENT:

Metalaxyl: *N*-(2,6-dimethylphenyl)-*N*-(methoxyacetyl) alanine methyl ester. 28.35%

INERT INGREDIENTS: 71.65%

TOTAL: 100.00%

Contains 2.65 lbs. Metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL
PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure,

Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

Product of South Korea Formulated in the USA

EPA Reg. No. 71532-22-55146

EPA Est. No.

Manufactured for
NUFARM AMERICAS INC.

AGT Division

150 Harvester Drive
Burr Ridge, IL 60527

Net Contents: 2.5 Gallons



FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the Poison Control Center at 1-800-222-1222.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING/AVISO: Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Coveralls over short-sleeved shirt and short pants, chemical-resistant gloves such as nitrile, butyl, neoprene, and/or barrier laminate, and chemical-resistant footwear plus socks. Also, wear a chemical resistant apron when mixing or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If not such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

GROUNDWATER ADVISORY STATEMENT

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or Viton.
- Chemical resistant footwear plus socks

In California — Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale.

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with Sebring 2.65 ST fungicide. Do not use for feed, food or oil purposes. Store away from feeds and foodstuffs."

This product is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, combination of Captan and Vitavax® registered formulations are compatible. Do not use with other seed treatment products unless previous experience assures compatibility.

This product may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: This product is a systemic fungicide having a specific mode of action. This product could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Nufarm Americas Inc. cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible **Sebring 2.65 ST** resistant strains of fungi, which may occur. FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

SEED TREATMENT

SEED	INSTRUCTIONS FOR USE
Canola* Mustard Seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply Sebring 2.65 ST as a seed treatment at the rate of 0.25-0.5 fl. oz. per 100 lbs. of seed. Use Restrictions: Do not harvest mustard greens. Rape greens grown and harvested from Sebring 2.65 ST treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from Sebring 2.65 ST treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption. *Not registered for use in California.
Carrots*	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.7 fl. oz. per 100 lbs. of seed. *Not registered for use in California.

Continued

SEED TREATMENT (continued)

SEED	INSTRUCTIONS FOR USE
Cotton	For <i>Pythium</i> seed rot and damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Soybeans	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Spinach*	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. Do not apply this product to spinach, which is destined to be grown in the greenhouse. Containers or bags containing Sebring 2.65 ST treated spinach seed must be labeled with this restriction. *Not registered for use in California.
Peanuts*	For <i>Pythium</i> seed rot and damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California.
Sorghum	For control of <i>Pythium</i> damping-off apply 0.375-0.75 fl. oz. of this product per 100 lbs. of seed applied as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75-1.5 fl. oz. this product per 100 lbs. of seed on cultivars, which possess sorghum downy mildew resistance. To those cultivars, which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5-3.0 fl. oz. of this product per 100 lbs. of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with this product.
Sunflowers	For control of systemic downy mildew: Apply this product as a seed treatment at the rate of 1.5-3.0 fl. oz. per 100 lbs. of seed.
Beets, Garden Sugar beets	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.7 fl. oz. per 100 lbs. of seed.

Continued

SEED TREATMENT (continued)

SEED	INSTRUCTIONS FOR USE
Forage Legumes	<p>Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).</p> <p>For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.</p>
Grain Crops	<p>Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale.</p> <p>For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For systemic downy mildew control of sweet corn: Apply this product as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Rice*	<p>For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. If using Sebring 2.65 ST treated rice do not use metalaxyl or mefenoxam for control of <i>Pythium</i> in water-seeded rice.</p> <p>*Not registered for use in California.</p>
Seed and Pod Vegetables	<p>Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, spring beans, wax beans, and lentils.</p> <p>For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For early season <i>Phytophthora</i> control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply this product as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Cucumbers*	<p>For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed. *Not registered for use in California.</p>
Turf Grasses	<p>For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply this product as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Forage Grasses	<p>Grasses grown for hay, grazing or silage corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.</p> <p>For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p>

SEED TREATMENT-EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

SEED	INSTRUCTIONS FOR USE
Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.5-1.7 fl. oz. per 100 lbs. of seed (32-110 ml per 100 kg of seed).
Corn	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply this product as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Onions	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed.
Millet	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply this product as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.
Peas	For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply this product as a seed treatment at the rate of 3.5 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. Do not apply this product to potatoes.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. Do not apply this product to potatoes.

Continued

SEED TREATMENT-EXPORT USE (continued)

SEED	INSTRUCTIONS FOR USE
Sorghum	For <i>Pythium</i> damping-off control: Apply this product as a seed treatment at the rate of 0.75-1.5 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply this product as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.
Sunflowers	For systemic downy mildew control: Apply this product as a seed treatment at 9.0 fl. oz. per 100 lbs. of seed.

Seed treated in accordance with the export use pattern must be labeled as follows:

FOR EXPORT ONLY - NOT FOR DOMESTIC SALES OR USE

Directions For Use at Reduced Rates in Combination With Other Fungicides: This product is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry of mist-type commercial seed treatment equipment.

SEED	INSTRUCTIONS FOR USE
Corn (field), Wheat, Oats, Barley, Rye	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply this product as a commercial seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply <u>only</u> in combination with EPA registered rates of seed treatment fungicides.
Soybeans, Field beans, Kidney beans, Navy beans, Pinto beans, Lentils, Peas	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply this product as a commercial seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply <u>only</u> in combination with EPA registered rates of broad spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply this product as a commercial seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply <u>only</u> in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Peanuts*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply this product as a commercial seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply <u>only</u> in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.
Rice*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply this product as a commercial seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply <u>only</u> in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store at temperatures above 40° F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

RETURNABLE/REFILLABLE CONTAINERS: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED “AS IS” BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(070307)

Sebring is a trademark of Nufarm Americas Inc.

Vitavax is a registered trademark of Crompton Corporation.

NOTES

NOTES