

10% Moly Liquid

By WINFIELD

Guaranteed Analysis

Molybdenum..... 10.0%

Derived from: Sodium Molybdate

Store above 32° F

KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid getting in eyes or on the skin. Keep Container closed when not in use.

STORE ABOVE 32° F

If Stored below 32° F, warm and agitate prior to use.

CAUTION: MOLYBDENUM CAN BE TOXIC TO PLANTS. MAKE SURE YOUR SOIL INDICATES THE NEED FOR ADDITIONAL MOLYBDENUM PRIOR TO SEED TREATMENT OR FERTILIZER USE. THE APPLICATION OF FERTILIZING MATERIALS CONTAINING MOLYBDENUM (Mo) MAY RESULT IN FORAGE CROPS CONTAINING LEVELS OF MOLYBDENUM (Mo) WHICH ARE TOXIC TO RUMINANT ANIMALS.

RECOMMENDED RATES AS SEED TREATMENT

Apply **10% Moly Liquid** at the rate of 3.5 to 5 fluid ounces per 100 pounds of seed. This rate will result in the application of 0.248 – 0.371 dry ounces of actual molybdenum per 60 pounds of seed.

10% Moly Liquid is formulated as a ready to apply material for soybeans. It supplies 0.248 dry ounces of actual molybdenum per bushel when applied at 2 fluid ounces per bushel of soybeans. It is compatible with most fungicide seed treatments, jar test before using.

Do not use treated seed for food, feed or oil purposes.

Recommended Applications and Rates as a Foliar Fertilizer

Alfalfa (seed and forage): 2 to 5 fl oz per acre every 2 to 3 cuts. Water rate: 20 gallons per acre.

Beans: 2 to 5 fl oz per acre when plants are 2 to 6 inches tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 20 gallons per acre.

Canola: For a single application, 2 fl oz per acre at onset of stem extension. For moderate deficiency, 2 fl oz at 4 to 6 leaf stage and 2 fl oz per acre at onset of stem extension. An extra application can be made 10 to 14 days later for a severe deficiency. Avoid flowering. Water rate: 20 gallons per acre.

Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower): 2 fl oz per acre from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at 10 to 14 day intervals from the first application. Water rate: 20 gallons per acre.

Carrots: 2 fl oz per acre. 10 to 14 days after transplanting or when crop is 6 inches tall. Water rate: 20 gallons per acre.

Cucurbits (field grown): 2 fl oz per acre 1 to 2 weeks after germination and at 4 leaf stage. Water rate: 20 gallons per acre.

F1683

Distributed and Guaranteed By:

Winfield Solutions, LLC

PO Box 64589

St. Paul, Minnesota 55164-0589

1/0427/6

Weight Per Gallon: 9.9 lbs. (4.49 kg)

NET WEIGHT: 24.75 lbs. (11.23 kg.)

NET CONTENTS: 2.5 Gallons (9.464 Liters)

WINFIELD™

Lettuce (field grown): 2 to 5 fl oz per acre 10 to 14 days after transplanting or emergence. Repeat at 7 to 10 day intervals if necessary. Water rate: 50 gallons per acre.

Melons (field grown): 2 fl oz per acre 1 to 2 weeks after germination and at 4 leaf stage. Water rate: 20 gallons per acre.

Peas: 2 to 5 fl oz per acre when plants are 2 to 6 inches tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 20 gallons per acre.

Soybeans: 2 to 5 fl oz per acre when crop is 2 to 6 inches tall, repeated at 10 to 14 day intervals if necessary. Water rate: 20 gallons per acre.

Sugar Beet: 2 fl oz per acre at 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 20 gallons per acre.

Tomatoes (field grown): 2 fl oz per acre when crop is 6 inches tall. Water rate: 50 gallons per acre.

Soil Application and Fertilizer Impregnation:

Soil Test Rating	Rate of 10% Moly
Mild Deficiency	10-20 fl. oz/acre
Moderate Deficiency	20-30 fl. oz/acre
Severe Deficiency	30-40 fl. oz/acre

May be applied in liquid fertilizer or impregnated on dry fertilizer.

PROPOSITION 65 WARNING: WARNING: This product may contain chemicals known to the State of California to cause cancer, birth defects or other reproductive harm.

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.