



Turf Fertilizer 0-0-22

with Dimension® Turf Herbicide & Ronstar®

FEATURES & USE TIPS:

- Provides preemergent control or suppression of the following annual weeds: goosegrass, crabgrass, annual bluegrass, and foxtail.
- Labeled for use on established or sprigged bermudagrass, zoysiagrass and St. Augustinegrass.
- Utilizes Ronstar and Dimension herbicides to maximize performance benefits of each active ingredient.
- Uniformly sized granules minimize particle segregation, which contributes to more accurate distribution.
- The readily available potassium increases resistance to heavy traffic and other turf stresses.
- Small particle formula provides more particles per square inch for better herbicide distribution.

PRODUCT CODE:

AGCDMR5

Date: 111307

Supersedes: 101702

Label: J7

PRODUCT CHARACTERISTICS:

Product Type:

Combination fertilizer,
Dimension® &
Ronstar® Herbicides

Particle Size:

SGN: 125

Net Weight:

50 lbs.

Coverage:

10,800 sq. ft. @
High rate

PRODUCT SPECIFICATIONS:

K - as K₂O (22%): Potassium Sulfate

S (7.48%) : Combined Sulfur

Active Ingredients: 0.125% Dithiopyr
1.000% Oxadiaxon

Dimension is a registered trademark of Dow AgroSciences L.L.C.; Ronstar is a registered trademark of Bayer; Andersons Golf Products is a trademark of The Andersons, Inc.

*To order, contact your local
Andersons Golf Products
distributor.*

*For more information, contact
Andersons at our website:
www.andersonsgolfproducts.com
or call 1-800-225-2639*



Turf Fertilizer 0-0-22®

with DIMENSION™ TURF HERBICIDE

This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

For sale and use by professional applicators only. Not for sale or use by homeowners/consumers.

Not for Use on Home Lawns

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 100 pounds per acre (equal to 5 pound Dimension active ingredient per acre), and is prohibited from use in Nassau and Suffolk Counties.

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O)	22.00%
7.48% Combined Sulfur	
F292	
1.000% Dimension	
0.125% Cinidon (dyclon)	
38.875% Capteggrass (Aconitum sativus)	
100.000% Penstemon (lanceolatum)	
Plant nutrients derived from potassium sulfate.	
ACTIVE INGREDIENTS:	
Cinidon (dyclon)	
Dithiopyr	
Total	

(Underlined letter is first letter seen in ran code on end of bag).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Leads, applicators and other handlers must wear:
• Long sleeve shirt and long pants, and
• Shoes plus socks
• Chemical resistant gloves made of any waterproof material

KEEP OUT OF REACH OF CHILDREN CAUTION

Some components of this product may cause eye irritation

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Leads, applicators and other handlers must wear:
• Long sleeve shirt and long pants, and
• Shoes plus socks
• Chemical resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions or wash

ables exist use detergent and hot water. Keep and wash PPE separately from other laundry.

USER USE PRECAUTIONS

The early postemergence crabgrass control described below is only effective on crabgrass power seeding equipment (slit seeders) designed to give good seed to soil contact.

For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT allow product or drift to contact foliage of flowers, vegetables, shrubs, trees, or fruit.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until product has been thoroughly watered in and grass has dried.

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contacts, if present after the first 5 minutes, then continue rinsing Call a poison control center or doctor for treatment advice

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff

turfs. Postemergence crabgrass suppression will be obtained only when this product is applied easily to the lawn or turf. So the practical benefit of this product and crabgrass products it can provide is that a 2-4 week longer period of time to make applications and still suppress crabgrass. This product is not intended to control large areas of established crabgrass, and must be used prior to crabgrass (goosegrass) for its control.

APPLICATION FREQUENCY AND TIMING:

For season-long control, this product should be applied as a single application in the spring for crabgrass (goosegrass) and crabgrass control.

DO NOT apply more than 40 lbs. per acre (0.5 lbs. Dimension and 4.0 lbs. Ronstar Alurate) per application, and no more than 600 lbs. per acre (0.75 lbs. Dimension and 6.0 lbs. Ronstar Alurate) per year, except in areas where there is a heavy weed infestation. In areas of heavy weed infestation, do not apply more than 800 lbs. of product per acre (1.0 lbs. Dimension and 8.0 lbs. Ronstar Alurate) per year.

Spring Applications: For single applications made in the spring, this product should be applied at the appropriate rate corresponding to one of the live control programs listed in table 1 below, depending on the user's location, the turfgrass growing height, and whether the use is considered to be pre-emergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use the Low Rate Program for preemergence control at sites where the turfgrass is cut high (e.g., rough), and/or (b) turf management or weed control has not been conducted during the previous year. This program provides 3-5 months of preemergence crabgrass (goosegrass) and crabgrass control, and so is especially useful in the transition zone, South of Orlando, FL, and/or in areas that do not receive a least one frost which naturally controls crabgrass (goosegrass) and crabgrass may require the use of a postemergence control product to eliminate existing weeds. This program provides early postemergence crabgrass suppression at sites where turfgrass is cut low (e.g., golf courses).

Use the High Rate Program for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and/or (b) turf management or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass (goosegrass) and crabgrass control, and so is especially useful in the southern and coastal south turf zones. South of Orlando, FL, and/or in areas that do not receive a least one frost which naturally controls crabgrass (goosegrass) and crabgrass may require the use of a postemergence control product to eliminate existing weeds. This program provides early postemergence crabgrass suppression at sites where turfgrass is cut low (e.g., golf courses).

USE & PRECAUTIONS

Do not apply to wet turf as discoloration may occur. Do not allow contamination of water used for irrigation and domestic use.

TURFGRASS SAFETY: This product may be used on lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand and have received at least two mowings following the seeding, sodding, or sowing before use. This product is not applicable to the weeds, which aids in providing excellent weed control while minimizing adverse effects to the turf. This product will not harm nearby established ornamentals when used according to label directions (avoid foliar contact).

WARM-SEASON GRASSES: This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to tolerate applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Area Classifications:	Low Rate	High Rate
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY & RI.	Ronstar: 1.5 lbs. ai per acre	Ronstar: 2.0 lbs. ai per acre
South: AL, AR, AZ, CA, GA, IA, MS, NC, NM, NV, OK, SC, TN & TX.	Dimension: 0.1875 lbs. ai per acre	Dimension: 0.25 lbs. ai per acre
Coastal South: HI, FL & southern coastal areas of AL, GA, LA, MS, NC, SC & TY.		

This product may cause temporary discoloration of Bermudagrass and St. Augustinegrass. This is normally outgrown within 2-3 weeks.

TIPS FOR IMPROVED CONTROL: For best results, apply this product within a few days after mowing and/or reseeding again or for a few days after the application. When treated lawns or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

SPRING APPLICATIONS: When used as directed to crabgrass (goosegrass) and crabgrass control in the spring, this product will also control the following weeds when applied prior to their emergence:

annual bluegrass	total (yellow & green)*
bamboo grass	Florida purple
Captegrass	herb
St. Augustinegrass	oxalis (butcher's broom) creeping & pineappaweed
Zoysia grass	purplegrass

For best results, water in the product as soon as practical after application. Thorough watering assures more even distribution of the herbicide and avoids gaps in coverage.

Avoid this product if spreader that would apply this product in narrow rows or concentrated bands. Before each application calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SGN 125

Net Wt. 50 Lbs. (22.68 kg)

SGN 120

Net Wt. 50 Lbs. (22.68 kg)

AGCDMR5
J7 (AGCBL KPRBN 1B)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

The suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 14,000 sq. ft. (0.33 acres) at 150 lbs. product - LOW RATE (0.45 lbs. 100 sq. ft.)

50 lbs. covers 10,800 sq. ft. (0.25 acres) at 200 lbs. product - HIGH RATE (0.6 lbs. 100 sq. ft.)

<p