

Turf Fertilizer with Millennium Ultra™ Herbicide 16-0-5



FEATURES & USE TIPS:

- Features Millennium Ultra™, a proven combination of 2,4-D, clopyralid and dicamba.
- 2,4-D provides broad-spectrum, post-emergent weed control, particularly on dandelions and other listed broad-leaf weeds.
- Clopyralid provides superior control of clovers.
- Dicamba provides added control
- 25% of total nitrogen is from NutriSphere-N® Nitrogen.
- SGN 150 for use on close-cut turf

PRODUCT SPECIFICATIONS:

N - (16%): Urea Nitrogen

K - as K₂O - (5%): Potassium Chloride

Cl - (4.24%): Chlorine Max

Active Ingredients: Dimethylamine Salt of 2,4-D (0.690%)
Monoethanolamine Salt of Clopyralid (0.094%)
Dimethylamine Salt of Dicamba (0.086%)

PRODUCT CODE:

AGC16NML4

Date: 071409
Supersedes: 000000
Label: E9

PRODUCT CHARACTERISTICS:

Product Type:

Combinatin fertilizer
& Millennium™ Ultra
Herbicide

Nitrogen Type:

NutriSphere-N™

Particle Size:

SGN: 150

Net Weight:

40 lbs.

Bags per pallet:

50

Coverage:

10,00 sq. ft.
@ 174 lbs/acre rate

Andersons Golf Products is a registered trademark of The Andersons, Inc.; Millennium Ultra is a trademark of Nufarm Americas Inc.; NutriSphere is a registered trademark of Specialty Fertilizer Products, LLC.

*To order, contact your local
Andersons Golf Products
distributor.*

*For more information, contact
Andersons at our website:
www.andersonsgolfproducts.com
or call 1-800-225-2639.*

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.

PRODUCT SHEET



Turf Fertilizer with

Millennium Ultra™ Herbicide 16-0-5

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE
AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE
SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH
FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, CLOPYRALID AND DICAMBA

- Formulated for fast herbicide response
- Contains NutriSphere®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-D*	0.930%
Clopyralid Acid**	0.071%
Dicamba Acid***	0.071%
OTHER INGREDIENTS:	
99.168%	
Total	100.000%

By Isomer Specific Method, Equivalent to:

* 2,4-D Acid	0.573%
** Clopyralid Acid	0.071%
*** Dicamba Acid	0.071%

EPA Reg. No. 228-343-9198
EPA Est. No. 9198-OH-TM, 9198-OH-2B, 9198-AL-001A
Undefined letter is first letter used in run code on bag

GUARANTEED ANALYSIS

Total Nitrogen (N)	16.00%
16.00% Urea Nitrogen†	
Soluble Potash (K ₂ O)	5.00%
Plant nutrients derived from urea and potassium chloride.	
Chlorine (Cl) Max.	4.24%
F282	

IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE SOLD TO AND USED BY
PROFESSIONAL APPLICATORS AND IS PROHIBITED FROM SALE AND USE IN MASSA-
CHUSETTS AND SUFFOLK COUNTIES.
IN CALIFORNIA, OREGON, AND WASHINGTON, TURFGRASS AND LAWN USES ARE
RESTRICTED TO GOLF COURSES ONLY.

This product cannot be used on Agricultural, Forest sites or Right-of-way.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, call (877) 325-1840

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

All handlers, applicators, and other handlers must wear:

- long-sleeved shirt and long pants
- steez plus socks
- chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for wash-
ables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing if pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	HOT LINE NUMBER
- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.	Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact: 1-877-325-6401 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ
ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH
LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Not for use on turf grown for sale or other commercial use as sod, or for commercial seed
production, or for research purposes.

This product is for use on Ornamental Turf Lawns (Industrial and Institutional), Parks, Cemeteries,
Athletic and Recreational Sports Fields, Golf Courses (Fairways, Aprons, Tees and Roughs) and
similar turf areas. *Excluding Bentgrass Tees.

The maximum number of broadcast applications per treatment site is 2 per year.

Do not apply this product in a way that will contact workers or other persons, either directly or
through drift. Only pre-treated handlers may be in the area during application. Do not allow people
(other than applicator) or pets on treatment area during application. Do not enter treatment areas
until dust has settled.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable
clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs,
trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

Do not use on residential turf. Do not collect grass clippings for mulch or compost. Do not send
clippings to a compost facility. Applicator must give notice to landowners/property managers if not
using grass clippings for mulching or composting.

WEEDS CONTROLLED

This product is designed to promote lush green growth in lawns and control many broadleaf weeds
such as: Alder, Annual yellow sweet clover, Amaranth, Aster, Austrian heliotropis, Bedstraw,
Barnyard grass, Broom, Broadleaf, Bird vetch, Bittersweet, Elder, wintercress, Black-eyed Susan, Black
Medic, Black Mustard, Black-seed plantain, Blessed thistle, Blue lettuce, Blue vernalis, Blue elder,
Bracted plantain, Brass buttons, Bristly oxeye, Broadleaf dock, Broadleaf plantain, Broomeed,
Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle,
Bur ragweed, Burnweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed,
bedstraw, Calceol, Calceol, Camp, Chickweed, Chiron, Chiron, Clover, Cockle, Cocklebur, Coffeebean,

Coffweed, Common chickweed, Common mullen, Common whitefish, Corn Chamomile,
Creeping jenny, Crimson clover, Cotton, Cuckweed, Cutly ridge, Dandelion, Dead
nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion,
False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flax bane (daisy), Flaxweed,
Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatshead,
Goldenrod, Ground ivy, Gumweed, Hairy bittersweet, Hairy fleabane, Hawkweed, Healed, Heartleaf
dymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cross, Hoary plantain, Hoary
vervain, Horeguslike, Hop clover, Horeguslike, Horegus, Indiana mallow, Ironweed, Jewelweed,
Jeweled, Knotweed, Knotweed, Lambsquarters, Lespedeza, Licorice, Lonicera, Malva,
Marshmallow, Matweed, Mexicanweed, Milk vetch, Milkweed, Milkweed, Mugwort, Morning
glory, Mousser chameleon, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle,
Orange hawkweed, Oxalis, Oxye daisy, Parsley-pert, Parsnip, Peatwort, Pennywort, Pennywort,
Peppercorn, Peppercorn, Pigeon, Pigeonweed, Pigeonweed, Pigeonweed, Pigeonweed, Pigeonweed,
henbit, Poison ivy, Poison oak, Pokeweed, Poorwill, Povertyweed, Picky leaf, Picky
sida, Primrose, Prostrate knotweed, Prostrate spurge, Prostrate spurge, Prostrate spurge,
Puncture vine, Purslane, Ragweed, Red clover, Redtop pigweed, Red sorrel, Redstem filaree,
Rhinoceros beetle, Rindweed, Roundleafed mallow, Rush, Russian pigweed, Russian
thistle, St. Johnswort, Serotinal primrose, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender
plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed,
Southern wild rose, Sowthistle, Spanshield, Spatterdock, Speedwell, Spiny Anemone, Spiny
cockbur, Spotted catnip, Spotted knapweed, Spotted spurge, Spurge, Spurred, String
nettle, Stinkweed, Stinkwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall
vervain, Tall spurge, Tansey root, Tansey, Tansey, Tansey, Tansey, Tansey, Tansey, Tansey,
Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica,
Vernon, Vetch, Virginia butterweed, Virginia creeper, Virginia pepperweed, Wayfared thistle,
Western dematis, Western salicy, White clover, White mustard, Wild aster, Wild buckwheat, Wild
card, Wild bulb-dock, Wild garlic, Wild geranium, Wild lettuce, Wild margot, Wild mustard,
Wild onion, Wild persim, Wild radish, Wild rose, Wild strawberry, Wild sweet potato, Wild vetch,
Willow, Whipweed, Woolly cotton, Woolly morniglovy, Woolly plantain, Woodsore, Wormseed,
Yarrow, Yellow pocket, Yellowflower pepperweed, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall.
Avoid contact with desirable shrubs, plants, vegetables, or flowers, especially when applying with a
rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply
to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2
days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3)
Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application.
Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48
hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn
is heavily overrun with weeds or if adverse low mowing conditions prevail and weeds are in a state of
poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to listed Warm Season Grasses - under conditions of inadequate moisture,
high temperature and high humidity, over-application may cause minor transitory discoloration. It
is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid
possible dehydration or browning of grass.

HOW TO APPLY: This 40-pound bag should be used on 10,000 square feet. Apply evenly with a
lawn spreader as a one pass application. A two pass application, cross-crossing the area at half
the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate
of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce
setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet
and adjust to apply at the rate of 1 pound per 250 square feet.

SPREADER SETTINGS			
Suggested spreader settings are not intended to replace calibration. Please calibrate your spreader prior to applying product. 40 lbs. treats 10,000 sq. ft. at the 174 lbs/acre rate			
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS 4.0 lbs/1,000 Sq. Ft.
Lb. active ingredient/acre (2,4-D; clopyralid; dicamba)			
1.20 0.12 0.12			
K			
Andersons AcoPro 2000, SP-2000 (cone 6)			
3 mph			
8 feet			
K			
Scotts R-8A (cone 6)			
3 mph			
8 feet			
J 1/2			
Lely Models			
WFR, WFR, HR, 1250			
26 feet			
4.3/4			
(PTD at 450 rpm) Pattern II-C			
Vicon (03 series)			
4.5 mph			
28 feet			
25			

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain
fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry
thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing
season. When used as directed, this product will supplement your regular fertilizer program by
providing 0.72 lb. of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please
consult your local extension office for questions regarding a regular fertilizer program.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured ware-
house or storage building. Do not stack more than two pallets high. Do not store near open
containers of fertilizer, seed or other pesticides.

PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has
leaked, clean up all spilled material. Improper disposal of excess pesticides, spray mixtures,
or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes
cannot be disposed of by use according to label instructions, contact your State Pesticide or
Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA
Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of
empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by
burning. If burned, stay away from smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT
CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE
FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND
SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY
KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE,
STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT
NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR
ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USE. UNLIMITED
CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT
BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS
USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION,
INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER
OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE
REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER,
AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO
MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO
THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL
MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, SPECIAL, INCIDENTAL, OR
CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES
RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND
DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER
OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION
TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS
OF MONEY CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON
OF RISKING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS
OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER
TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS
CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYERS OR USERS
EXCLUSIVE REMEDY, AND MANUFACTURERS OR SELLERS' TOTAL LIABILITY SHALL BE
FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaim-
ers, or limitations on liability, do not use the product, and return it unopened to the Seller,
and the purchase price will be refunded.

Distributed by:

The Andersons Lawn Fertilizer Division, Inc.
P.O. Box 119

Maumee, Ohio 43537

www.andersonsgolproducts.com

SGN 150
Net Wt. 40 Lbs. (18.14 kg)



AGC16NML4

E9 AGCBLK(PBFIN.1B)

0 318651 313931 6

The Andersons logo is a registered trademark of The Andersons, Inc.; Millennium Ultra is a
trademark of Nuliam Americas Inc.; NutriSphere is a registered trademark of Specialty Fertilizer
Products, LLC.