

SFT™ Fertilizer with 1% Ronstar® Herbicide

22-0-8



FEATURES & USE TIPS:

- Provides preemergent control or suppression of the following annual weeds: goosegrass, crabgrass, *Poa annua*, and foxtail.
- Utilizes SFT™ Formulation Technology, which will reduce the potential for turf discoloration and injury compared to conventional fertilizers and Ronstar products.
- Labeled for use on established or sprigged bermudagrass, zoysiagrass, St. Augustinegrass and Seashore paspalum* (see precautions on label).
- Utilizes Ronstar herbicide which can be used prior to sprigging warm season turf without “tacking down” concerns.
- Small particle (SGN 150) sizing for use on low cut fairways.
- The 22-0-8 formulation contains Poly-S® slow release nitrogen, to provide prolonged feeding action with minimum surge growth.

*Not registered in California.

PRODUCT SPECIFICATIONS:

N - (22.00%): Urea Nitrogen*
*22.00% Slowly available nitrogen from polymer coated sulfur coated urea

K - as K₂O (8.00%): Potassium sulfate

S - (4.76%): Total Sulfur
2.04% Free Sulfur
2.72% Combined Sulfur

Active Ingredient: Oxadiazon (1.00%)

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.

PRODUCT CODE:

AGC222R50

Date: 011409

Supersedes: 101207

Label: K8

PRODUCT CHARACTERISTICS:

Product Type:

SFT™ Fertilizer &
Ronstar® Herbicide

Nitrogen Type:

Poly-S®

Particle Size:

SGN: 150

Net Weight:

50 lbs.

Coverage:

10,800 sq. ft. @
medium rate

Ronstar® is a registered trademark of Bayer.; SFT is a trademark of The Andersons, Inc.; Andersons Golf Products® is a registered trademark of The Andersons, Inc.

*To order, contact your local
Andersons Golf Products
distributor.*

*For more information, contact
Andersons at our website:
www.andersonsgolfproducts.com
or call 1-800-225-2639*



SFT™ Fertilizer with 1% RONSTAR® Herbicide

For Sale to, Use and Storage by Professional Turf Maintenance Personnel only.
Not for sale to or use by homeowners/consumers.
Not for Use On Home Lawns

22-0-8

- Contains Ronstar® herbicide, which provides preemergent control or suppression of crabgrass, *Poa annua*, goosegrass and foxtail during Bermudagrass springing.
- Utilizes SFT™ technology

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Loaders, applicators and other handlers must wear:
- Long-sleeved shirt, and long pants, and
 - Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY/RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed

Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8851 for emergency medical treatment information.

Environmental Hazards: This pesticide is toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

ACTIVE INGREDIENT

Oxadiazon†	1.0%
OTHER INGREDIENTS	99.0%
Total	100.0%

†CAS Number 16866-30-9

EPA Reg. No. 9198-0203

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in min. runs on ban

GUARANTEED ANALYSIS

Total Nitrogen (N).....	22.00%
22.00% Urea Nitrogen*	
Soluble Phosph (P ₂ O ₅).....	8.00%
Sulfur (S).....	4.76%
2.04% Free Sulfur (S)	
2.72% Combined Sulfur (S)	

Plant nutrients derived from potassium sulfate and polymer coated sulfur coated urea.
*22.00% Slowly available nitrogen from polymer coated sulfur coated urea

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coversalls, shoes plus socks, and chemical resistant gloves made of any waterproof material.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, sod farms or greenhouses.

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated areas (even those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

TURF USES

For use on established or sprigged Bermudagrass, Zoysiagrass, Seashore Paspalum* (see precautions), Tropic Lab (paspalum hybridum), St. Augustinegrass, perennial Bluegrass, Perennial Ryegrass and Tall Fescue turf, such as lawns, parks, golf courses and sod farms. Use for preemergent control or suppression of sprouting.

Weed Grasses:

Crabgrass
Goosegrass
Annual Sage*

Broadleaf Weeds:
Oxalis
Florida Pusley
Virginia Buttonweed

*Not currently registered in California

Yellow foxtail
Green foxtail
Annual Bluegrass (Poa annua)
Field Sandbur*

Barnyardgrass
Cenchrus
Common Puncture
Pigweed
Veronica Speedwell
Common Groundsel

How to Use

- Apply prior to or after springing. 150 to 200 lbs. of product per acre (1.5 to 2.0 lbs. Alacre) for suppression or control of listed annual weeds in bermudagrass, zoysiagrass, and St. Augustinegrass.
- Apply 200 to 300 lbs. of product per acre (2.0 to 3.0 lbs. Alacre) for control of listed annual weeds during bermudagrass, zoysiagrass, seashore paspalum and tropic lab (paspalum hybridum) establishment from spring.
- After application of this product on newly sprigged bermudagrass, zoysiagrass, and St. Augustinegrass, wait until the grass has filled in and the stores have rooted down before application of other herbicides, or growth regulator products.
- On Seashore paspalum make application 10-14 days after springing.
- For established turf, apply at 200 to 300 lbs. of product per acre (2.0 to 3.0 lbs. Alacre).

When to Use

Use this product to control or suppress the listed annual weeds when establishing new turf areas or when treating established turf that has been thinned from winter injury, insects, diseases or other factors.

- Bermudagrass - make application immediately prior to or immediately after springing.
- Zoysiagrass and St. Augustinegrass - make application prior to or after springing.
- Seashore Paspalum - make application 10 to 14 days after springing. Applications made close to the time of springing may cause objectionable Seashore Paspalum injury.
- Tropic Lab - application must be made 0-12 days after springing. Do not use in areas where endangered Hawaiian Goose or nene (*Brania sandwicensis*), Hawaiian duck (*Anas wyvilliana*), Hawaiian Stilt (*Himantopus mexicanus knudseni*) and Hawaiian Monkseal (*Calidris choropus sandwichensis*) may feed.

For best results, water-in the product as soon as practical after application.

*Not currently registered in California

PRECAUTIONS

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

Avoid contact with flowers and shrubs.

Do not apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment.

This product should not be used on putting greens or tees because of the varied cultural practices used.

Do not apply to Bermgrass turf mowed less than 3/8 inch.

Do not apply to wet turf as discoloration may occur.

For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 240 lbs. per acre (2.4 pounds al acre).

This product may cause temporary discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This is normally out-grown within 2-3 weeks.

On Seashore Paspalum wait 10-14 days after springing before application. An application made close to the time of springing may cause objectionable injury.

Do not apply to Dichondra or Centipedegrass.

Do not apply more than 300 lbs. of product per acre (3.0 lbs. Alacre) per single application.

Do not apply more than 600 lbs. of product per acre (6.0 lbs. Alacre) per year except in areas where there is heavy weed infestation. If areas of heavy weed infestation, do not apply more than 800 lbs. of product per acre (8.0 lbs. Alacre) per year.

Disburse of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. **PESTICIDE STORAGE:** Store in clean, dry place. Re-seal opened bag by tucking top down and securing. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess product, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. To the extent consistent with applicable law, by using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law, manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. It is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/ JURISDICTION TO STATE/ JURISDICTION. (Rvw070705)

SUGGESTED SPREADER SETTINGS				
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.				
50 lbs. treats 10,800 sq. ft. (0.25 acre) at the 200 lbs. product/acre - MEDIUM RATE (4.59 lbs/1,000 sq. ft.)				
50 lbs. treats 8,700 sq. ft. (0.20 acre) at the 250 lbs. product/acre - HIGH RATE (5.74 lbs/1,000 sq. ft.)				
50 lbs. treats 7,200 sq. ft. (0.17 acre) at the 300 lbs. product/acre - HEAVY RATE (6.89 lbs/1,000 sq. ft.)				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			MEDIUM RATE 20 lbs. al/acre	HEAVY RATE 3.0 lbs. al/acre
Andersons AccuPro 2000, SP2000 (cone 8)	3 mph	12 feet	L 1/4	M 1/4
Scotts R-8A (cone 8)	3 mph	11 feet	K 3/4	M 3/4
Lely Models WTR, WFR, HR, (250 PTO at 450 rpm) Pattern 1	4.5 mph	36 feet	7 1/2	8 1/2
Vicon O3 Series	4.5 mph	30 feet	30	34
				39

IMPORTANT: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

ATTENTION: This product contains a chemical known to the State of California to cause cancer.


Manufactured by:

The Andersons Lawn Fertilizer Division,
PO Box 119, Maumee, OH 43537
www.andersonsgolfproducts.com

Andersons Golf Products is a registered trademark of The Andersons, Inc.; SFT is a trademark of The Andersons, Inc.; Ronstar is a registered trademark of Bajet.

SGN 150

Net Wt 50 Lbs (22.68 kg)



AGC222R50

K8® (AGCBLKPRN-1B)