

BEST BALE®

A HAY TREATMENT PRODUCT

ACTIVE INGREDIENTS:

Crude Protein	17.00%
From non-protein (ammoniacal) nitrogen	
Total Phosphorus	2.00%
Potassium (K), Sulfur (S), Calcium (Ca), Magnesium (Mg), Chloride (Cl), Sodium (Na), Iron (Fe), Manganese (Mn), Zinc (Zn), Boron (B), Copper (Cu), and other minerals	<1.00%
Naphthaleneacetic acid (NAA)	0.000001%
Indole-3-butyric acid (IBA)	0.0000015%
Cellulase	0.36%
Lactobacillus acidophilus	1.1x10.3 CFU/ml
Organic acids including HUMIC and FULVIC	2.0%

Derived From Urea, Iron Chelate, Zinc Chelate, Manganese Chelate, Polyacrylamide Sodium Molybdate, Cobalt Chloride, NAA, IBA, Cellulase, Lactobacillus acidophilus
Chelating agents are Hydroxyethylethylenediaminetriacetate and Ethylenediaminetriacetate.
Also contains 14.0% organic acid complex derived from leonardite and/or composted manure.

What To Do In Case Of Contact:

If swallowed: Immediately call a physician. Drink large amounts of water. Induce vomiting. **If in eyes:** Flush with large amounts of water for 15 minutes. Get medical attention.

If on skin: Wash exposed area with water.

General Information:

BEST-BALE® is a hay treatment product applied pre-baling where spoilage due to rain or high moisture conditions is a concern.

DIRECTIONS FOR USE:

• **BEST-BALE®** should be incorporated with water as a carrier into swathed hay at the header or crimper/roller. **BEST-BALE®** could also be applied directly to the windrowed hay or at the baler, but this application would be less desirable.

RATE OF APPLICATION:

HAY: 4-6 ounces of **BEST-BALE®** with 2-5 gallons of water per ton of dry hay (15% moisture). **SILAGE:** 20-30 ounces of **BEST-BALE®** with 5-10 gallons of water per ton (75% moisture).

STORAGE AND DISPOSAL:

Keep product in original container. Do not transfer into food or drink containers. Triple rinse container when empty for recycling. Always dispose of container in accordance with local, state and/or federal regulations.

OTHER NOTES:

- These recommended rates and methods of application are basic guidelines intended to simplify product usage and maximize results. There may be specific conditions which would require adjustments.
- Consult your local HUMA GRO® representative or other agricultural specialist for specific recommendations.

WARNING: Application of this material in excess may result in forage crops containing levels of Molybdenum (Mo) which are toxic to ruminant animals.

Conditions of sale:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

CAUTION: KEEP OUT OF REACH OF CHILDREN



☐ **2.5 Gallons • Net Wt. 23.0 lbs.**
9.46 Liters • Net Wt. 10.43 Kilograms

☐ **5 Gallons • Net Wt. 56.0 lbs.**
18.93 Liters • Net Wt. 20.86 Kilograms

● 9.20 lbs. per gallon @ 68° F.
1.10 Kilograms per Liter @ 20° C.
pH 7.9