BORON

GUARANTEED ANALYSIS

Derived from: Boric Acid

METHOD OF APPLICATION	SUGGESTED RATE PER ACRE	
	Field Crops	Tree or Vine Crops
Foliar band application at 50% coverage	up to 1 pint	
Foliar broadcast or sprinklers: solid, set, linear, pivot (100% speed)	up to 1 quart	up to 2 quarts
Soil banded or injected through drip tape or micro sprinklers	up to 1 quart	up to 2 quarts
Soil broadcast spray incorporated, flood or furrow irrigated	up to 2 quarts	up to 1 gallon

DIRECTIONS FOR USE

•Shake well before use. •Designed to be applied by both soil and/or foliar application. •BORON should be applied when tissue and soil analysis identifies the need and proper rate of application for this product. •Best results will be obtained when application is concentrated in the active root zone or on the leaf surface. •Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. •If compatibility is in question, jar test a small quantity. •Apply foliar sprays with sufficient water to ensure uniform coverage without running off leaf surfaces. •Do not apply this product foliarly in concentrations greater than 5%. •For maximum results, apply in the early morning or in the evening when air temperatures are optimum for foliar application. •Do not apply HUMA GRO® nutrients during the hottest time of the day or to plants under drought or flood stress. •Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop. •Applications can be made as often as every 7 to 10 days as needed. •Consult your local HUMA GRO® Representative or other agricultural specialist for crop-specific recommendations. •Do not store this product below 50°F (10°C) or above 90°F (30°C).

This product is intended as a supplement to a complete fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.

Conditions of sale:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

CAUTION: KEEP OUT OF REACH OF CHILDREN

WARNINGS: This product contains Boron (B), which may be injurious to certain crops. The use of this fertilizing material on any crop(s) other than those recommended may result in serious injury to the crop(s).

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm



Produced by
Bio Huma Netics, Inc. ©2005-2012
1331 W. Houston Ave, Gilbert, AZ 85233
480-961-1220 • www.humagro.com

9.87 lbs. per gallon @ 68° F.
 1.18 Kilograms per Liter @ 20° C.
 pH 6.5 - 7.5

1 Liter • Net Wt. 2.61 lbs.
1 Liter • Net Wt. 1.18 Kilograms

4 Liters • Net Wt. 1.18 Kilograms
4 Liters • Net Wt. 4.73 Kilogram

4 Liters • Net Wt. 4.73 Kilograms

2.5 Gallons • Net Wt. 24.68 lbs.

9.46 Liters • Net Wt. 11.19 Kilograms

5 Gallons • Net Wt. 49.35 lbs.
18.93 Liters • Net Wt. 22.38 Kilograms

275 Gallons • Net Wt. 2,714.25 lbs. 1,041.15 Liters • Net Wt. 1,230.95 Kilograms



HG-020-09132012