



BeanOil®

Spray Tank Adjuvant For Pesticides

PRINCIPAL FUNCTIONING AGENTS:

Soybean oil	60.0%
Alkylphenol ethoxylate, alcohol ethoxylate	40.0%
TOTAL:	100.0%

Ingredients exempt from tolerance under 40 CFR 180.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See **FIRST AID** Below

Read Entire Label Before Using This Product

Net Content: _____

FIRST AID
<p>IF IN EYES:</p> <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>IF ON SKIN OR CLOTHING:</p> <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>IF SWALLOWED:</p> <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>IF INHALED:</p> <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes eye and skin irritation. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, smoking tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Wear chemical-resistant gloves, long-sleeved shirt and long pants, and shoes plus socks.

ENVIRONMENTAL HAZARDS

Do not contaminate water sources by cleaning of equipment or disposal of wash water.

USE INFORMATION

BEANOIL is a superior blend of special emulsifier-surfactants and highly refined soybean oil that is designed for use with a broad range of pesticides where an oil concentrate adjuvant is recommended. This product may be used at rates of 0.25 to 0.75% v/v with herbicides, insecticides, fungicides, fertilizers and micronutrients. The addition of this product to a spray tank improves pesticide applications by modifying wetting and contact activity of the spray, enhancing penetration and providing uniform deposition of spray deposit to the leaf surface. This product contains anti-foam to suppress foam development in the tank mix combinations.

DIRECTIONS FOR USE

This product may be used with products registered for: Agricultural, Industrial, Forestry Site Prep, Municipal, Non-cropland, Utility and Roadside rights-of-way, Ornamentals, Turf and other uses. The addition of an adjuvant to some pesticides or tank mix combinations may cause phytotoxicity to foliage or fruit of susceptible crops.

Refer to the label of the tank mixture partner(s) for any additional use instructions or restrictions. Always follow the most restrictive label.

This product may be applied by high or low volume ground, aerial or any type of equipment recommended by pesticide labels.

Not for aquatic use.

MIXING: To prepare sprays containing this product, fill the spray tank half full with water, add the proper amount of pesticide, fertilizer or micronutrient as directed by its label and then add this product. Maintain agitation during the entire process. Continue agitation before and during each application to ensure a uniform spray.

Note: Care should be exercised when spray containing this product, is applied to new crop varieties or highly inbred crop lines not previously sprayed with this product. It is recommended that a small test area be treated before spraying large areas. Use care when pesticides are more active, such as during periods of hot or humid conditions.



Manufactured By:

Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

Crop	Problem/Situation	Pesticide*	Rate of This Product
Citrus	Annual and Perennial grass weeds	Fusilade®, Gramoxone®, Poast®, Prism®	0.25 to 0.50% v/v or 8 to 16 fl. ozs. per acre
Corn (Field, Seed, Silage, Sweet)	Annual and Perennial grasses, Broadleaf weeds	Accent®, Atra-5™, Atrazine 4L or 90DF, Banvel®, Basagran®, Basis Gold®, Clarity®, Guardsman®, Resource®	0.50 to 0.75% v/v or 8 to 16 fl. ozs. per acre
Cotton	Annual and Perennial grasses, Broadleaf weeds	Cobra®, Diuron 4L & 80 Herbicide, Fusilade, Poast, Poast Plus, Select®, Staple®	0.25 to 0.50% v/v or 8 to 16 fl. ozs. per acre
Cotton	Post-emergent grass (Spot or small area treatment)	Fusilade, Poast	8 to 16 fl. ozs. per 25 gals.
Cotton	Boll openers	Def® 6, Harvade®	8 to 16 fl. ozs. per acre
Cotton	Defoliation	Defol® 5, Defol 750, Def 6, Harvade	8 to 16 fl. ozs. per acre
Peanuts	Annual and Perennial grasses, Broadleaf weeds	Basagran, Poast, Pursuit®, Cobra, 2,4-DB	0.50 to 0.75% v/v or 8 to 16 fl. ozs. per acre
Peanuts	Ascochyta web blotch, Blackhull, Cercospora leaf spot, Rust	Benlate® with Manzate® (Do not use Benlate alone), Cobra, 2,4-DB	4 to 8 fl. ozs. per acre
Pecans	Brown leaf spot, Liverspot, Powdery mildew, Scab	Benlate Fungicide (Do not use Benlate alone)	4 to 8 fl. ozs. per acre
Pome and Stone fruits	Black rot, Fly speck, Fruit rots, Powdery mildew, Scab, Sooty blotch	Benlate Fungicide (Do not use Benlate alone)	4 to 8 fl. ozs. per acre
Potatoes	Annual and Perennial grasses	Gramoxone, Poast	0.25 to 0.50% v/v or 1 to 2 pts. per acre
Potatoes	Vine desiccation	Defol 5, Defol 750	8 to 16 fl. ozs. per acre
Rice	Annual and Perennial grass weeds	Basagran, Propanil	8 to 16 fl. ozs. per acre
Sorghum or Milo	Annual and Perennial grass weeds	Atrazine 4L or 90DF, Atra-5	8 to 16 fl. ozs. per acre
Soybeans	Annual and Perennial grass weeds	Basagran, Classic®, Cobra, Fusilade, Poast, Pursuit, Resource, Synchrony® STS	0.25 to 0.75% v/v or 8 to 16 fl. ozs. per acre
Soybeans	Post-emergent grass (Spot or small area treatment)	Fusilade, Poast	8 to 16 fl. ozs. per 25 gals.
Sugar beets	Annual and Perennial grass weeds	Poast, Prism	0.50 to 0.75% v/v or 1 pt. per 20 gals. of water
Sugarcane	Annual and Perennial grass weeds	Atrazine 4L or 90DF, Atra-5, Diuron 4L, Diuron 80 Herbicide	4 to 16 fl. ozs. per acre
Ornamental, Nursery, and other Non-food crops	Post-emergent grass	Poast	16 fl. ozs. per acre

* Follow pesticide label(s) for use rates and instructions.

NON-CROP USE		
Application Site	High-Volume Ground (per 100 gals.)	Low-Volume Ground or Aerial (per acre)
Conifer Release* (Herbaceous and Woody)		4 fl. ozs. Herbaceous 8 fl. ozs. Woody (Based on 10 gallons per acre total volume)
Ditchbank	1 to 4 pts.	2 to 4 pts.
Forestry Site Prep	1 to 4 pts.	2 to 4 pts.
Non-Crop Applications	1 to 4 pts.	2 to 4 pts.
Utility and Roadside Rights-of-way	1 to 4 pts.	2 to 4 pts.

* Do not exceed 0.25% by spray volume for Herbaceous conifer release or 0.75% by spray volume for Woody conifer release.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PRODUCT STORAGE: Store in a cool, dry, well-ventilated area. Avoid high temperatures. Store above freezing.

PRODUCT DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler, visit the ACRC web page at www.acrecycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.