

Drexel

# Carbaryl 80S

For Agricultural or Commercial Use Only

**ACTIVE INGREDIENT:**

Carbaryl 1-naphthyl  
N-methylcarbamate: ..... 80.0%

**OTHER INGREDIENTS:** ..... 20.0%

**TOTAL:** ..... 100.0%

## KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See **FIRST AID** Below

EPA Reg. No. 19713-50

EPA Est. No. 19713-GA-1

Net Contents: \_\_\_\_\_

### FIRST AID

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.

**IF INHALED:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**GENERAL**

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.** If poisoning is suspected in animals, contact a veterinarian.

**ANTIDOTE STATEMENT**

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended. See **NOTE TO PHYSICIAN.**

**NOTE TO PHYSICIAN:** Treat symptomatically. Overexposure to materials other than this product may have occurred. Carbaryl is an N-methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product has been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

(Continued)

**NOTE TO PHYSICIAN (Cont.)**

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent over-atropinization. Improve tissue oxygenation as much as possible before administering atropine to minimize the risk of ventricular fibrillation. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended. To aid in confirmation of diagnosis, urine sample should be obtained within 24 hours of exposure and immediately frozen.

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**

**WARNING:** MAY BE FATAL IF SWALLOWED. HARMFUL IF ABSORBED THROUGH SKIN, INHALED, OR IF IN EYES. Do not breathe vapors, dust or spray mist. Do not get in eyes, on skin or on clothing. Keep out of reach of children and domestic animals.

**OVEREXPOSURE MAY CAUSE:** Salivation, watery eyes, pinpoint eye pupils, blurred vision, muscle tremors, difficult breathing, excessive sweating, abdominal cramps, nausea, vomiting, diarrhea, weakness and headaches. **IN SEVERE CASES, CONVULSIONS, UNCONSCIOUSNESS AND RESPIRATORY FAILURE MAY OCCUR. SIGNS AND SYMPTOMS OCCUR RAPIDLY FOLLOWING OVEREXPOSURE TO THIS PRODUCT.**

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:** Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:** 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Discharge from Rice fields may kill aquatic and estuarine invertebrates. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

Manufactured By:

**Drexel Chemical Company**

P.O. BOX 13327, MEMPHIS, TN 38113-0327

**SINCE 1972**

• 50SP-0901++

CARBARYL 80S Page 1 of 11

**BEE CAUTION:** MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service for further information.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product. Strictly observe label directions and precautions. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 12 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, Neoprene Rubber, Polyvinyl Chloride (PVC) or Viton, shoes plus socks and chemical-resistant headgear for overhead exposure.

#### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

The area being treated must be vacated by unprotected persons. Keep unprotected persons out of treated areas until sprays have dried.

#### **GENERAL PRECAUTIONS AND RESTRICTIONS**

CARBARYL 80S readily disperses in water to form a spray which may be applied by air or ground.

**Plant Response Precautions:** Application to wet foliage or during periods of high humidity may cause injury to tender foliage.

Do not use on Boston ivy, Maidenhair fern and Virginia creeper as injury may result. Carbaryl may also injure Sand and Virginia pines. The use of adjuvants may increase the potential for crop injury to sensitive crops.

#### **PRE-HARVEST AND GRAZING RESTRICTIONS AND LIMITATIONS**

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of labeled crops bearing probable carbaryl residues when this product is used in accordance with the label directions. If used as directed, treated forage may be grazed or used as feed for dairy and meat animals without causing illegal residues in meat or milk. Do not apply at greater rates or at more frequent intervals than stated on the label. To do so may result in illegal residues in crops, meat and milk. Do not use reclaimed irrigation water from crops treated with carbaryl on crops for which carbaryl tolerances are not established. Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

#### **APPLICATION STATEMENTS**

Calibrate and adjust application equipment to ensure proper rate and accurate placement. To clean spray system after use, drain and flush with water and detergent mixture. Rinse thoroughly with clean water. Refer to the "STORAGE AND DISPOSAL" section for disposal instructions.

**Note:** Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Spray deposits on painted or stained surfaces or finishes (i.e., cars, houses, trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid applications to surfaces where visible spray residues are objectionable.

**RESISTANT SPECIES NOTICE:** All references to Armyworms on the crops listed on this label refer to the species, *Pseudaletia unipuncta*, often called the "True armyworm". Except where indicated otherwise, this product is not registered for the control of other Armyworm species. Regional differences have been noted in the susceptibility of certain strains of Colorado potato beetles, Diamondback moths, Fall armyworms and Southern green stink bugs to carbaryl. If local experience indicates inadequate control, use an alternative pesticide.

#### **MIXING, LOADING AND HANDLING INSTRUCTIONS**

TO ASSURE A UNIFORM SUSPENSION, AGITATE, STIR OR RECIRCULATE ALL CONTAINERS OF THIS PRODUCT PRIOR TO USE. Remove

oil, rust, scale, pesticide residues and other foreign matter from mix tanks and entire spray system. Flush with clean water. Fill spray or mix tank with ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of this product and then the remaining volume of water. Include rinse water from container. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURE FOR PROLONGED PERIODS OR DEGRADATION OF CARBARYL MAY OCCUR. Local water conditions may also accelerate the degradation of spray mixtures containing carbaryl. See "COMPATIBILITY STATEMENT" below.

#### **COMPATIBILITY STATEMENT**

This product, when diluted with at least an equal volume of water, is compatible with a wide range of pesticides. It is not compatible with diesel fuel, kerosene, fuel oil or aromatic solvents. If compatibility with another product and the resulting crop response is unknown, the mixture should be tested on a small scale. Curdling, precipitation, greasing, layer formation or increased viscosity are symptoms of incompatibility. Incompatibility will reduce insect control and may cause application and handling difficulties or plant injury. Observe all precautions and limitations on labeling of all products used in mixtures. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD THIS PRODUCT TO AT LEAST AN EQUAL VOLUME OF WATER, MIX THOROUGHLY AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE. DO NOT APPLY TANK MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS OR PLANT INJURY.

Carbaryl is unstable under highly alkaline conditions and mixture with strong bases, such as Bordeaux, lime-sulfur and casein-lime spreaders, will result in chemical degradation of the insecticide. Do not use this product in water with pH values above 8.0 unless a buffer is added. If necessary, water should be buffered to neutral (pH = 7.0) before adding this product to the spray tank. Overhead irrigation with alkaline or muddy water after application will also accelerate chemical degradation and may result in reduced insect control.

#### **APPLICATION PROCEDURES AND PRECAUTIONS**

On all crops, use sufficient gallonage to obtain thorough and uniform coverage. Observe crop label instructions for specific directions regarding spray volume where they occur. Calibrate spray equipment to deliver the required volume. Use of 50-mesh slotted strainers in spray system and 25-mesh slotted strainers behind nozzles is recommended.

**Ground Application:** Apply in sufficient volume for adequate coverage on all crops and sites. To prepare small volumes of spray mixture, use 6.5 grams (approximately 1.2 tablespoons) of this product in an adequate amount of water and apply to 500 square feet, where rates of 1¼ pounds per acre are indicated.

**Aerial Application:** For adequate distribution, use at least 10 gallons of spray mixture per acre for application for tree and orchard crops or at least 2 gallons of spray mixture per acre for application to other crops.

**Sprinkler Irrigation Systems:** Apply this product only through sprinkler irrigation systems including center pivot and solid set. Do not apply this product through any other type of irrigation system.

**Spray Preparation:** First prepare a suspension of this product in a mix tank. Fill tank with ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of this product and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label, per 1 to 4 gallons of water are recommended.) Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of this product into the irrigation waterline so as to deliver the desired rate per acre. The suspension of this product should be injected with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

**Note:** When treatment with this product has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

**General Precautions For Applications Through Sprinkler Irrigation Systems:** Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation

line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained. Crop injury, lack of effectiveness or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

#### SPECIFIC USE DIRECTIONS

**Crop/Site Groupings:** Asparagus, Brassica leafy vegetable crops, Cereal grain crops (Field and Popcorn, Grain sorghum, Rice, Sweet corn, Wheat and Proso millet); Cucurbit vegetables, Flax, Forage crops (Alfalfa, Birdsfoot trefoil, Clovers, Pasture and Grasses grown for seed, Rangeland), Fruiting vegetables, Leafy vegetables, Legume vegetables, Non-cropland (Conservation Reserve Program—Ditchbanks, Hedgerows, Right-of-ways, Roadsides, Wasteland), Okra, Peanuts, Prickly pear cactus, Root and Tuber crops, Sugar beets, Sweet potatoes, small Fruits and Berries, Sunflower, Tobacco, Tree fruit crops (Citrus fruits, Olives, Pome fruits, Stone fruits), Tree nut crops (Pistachios, Tree nuts), Forested areas and Rangeland trees, Trees and Ornamentals, Turfgrass, control of specific pests across multiple sites (adult Mosquito control, Grasshoppers, Imported fire ants, Ticks which vector lyme disease) and nuisance pests around buildings.

#### INSECT CONTROL

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, consultants or other qualified authorities to determine appropriate threshold levels for treatment and specific use information in your area. Where a dosage range is indicated, use the lower rate on light to moderate infestations, young plants and early instars and use the higher rate on heavy infestations, mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

#### ASPARAGUS

Crop	Pest	Lbs. of This Product Per Acre
Asparagus	Apache cicadas, Asparagus beetles, Cutworms	1¼ to 2½
	SPECIFIC DIRECTIONS: Repeat application as necessary up to a total of 3 times prior to harvest, or a total of 5 times per crop, but not more often than once every 3 days. For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
	Apache cicadas, Asparagus beetles, Cutworms	2½ to 5
	SPECIFIC DIRECTIONS: Application to ferns or brush growth following harvest of spears: Repeat applications as necessary but not more often than once every 7 days. Do not make more than a total of 5 applications per year to spears and ferns combined.	

**RESTRICTIONS AND PRECAUTIONS:** Do not apply within 1 day of harvest. Do not apply more than a total of 7½ lbs. per acre before harvest of spears. Do not apply more than a total of 12½ lbs. per acre per year.

#### BRASSICA LEAFY VEGETABLE CROPS

Repeat applications as needed up to a total of 4 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese cabbage, Collards, Kale, Kohlrabi, Mustard greens	Flea beetles, Harlequin bugs, Leafhoppers	¾ to 1¼
	Armyworms, Aster leafhoppers, Corn earworms, Diamondback moths, Fall armyworms, Imported cabbageworms, Lygus bugs, Spittlebugs, Stink bugs, Tarnished plant bugs	1¼ to 2½

**RESTRICTIONS AND PRECAUTIONS:** For Broccoli, Brussels sprouts, Cabbage, Cauliflower and Kohlrabi, do not apply within 3 days of harvest. For Chinese cabbage, Collards, Kale and Mustard greens, do not apply within 14 days of harvest. Do not apply more than a total of 7½ lbs. per acre per crop.

#### CEREAL GRAIN CROPS

Repeat applications as needed up to a total of 4 times but not more often than once every 14 days. Optimum timing and good coverage are essential for effective control.

Crop	Pest	Lbs. of This Product Per Acre
Field corn, Popcorn	Armyworms, Chinch bugs, Corn earworms, Corn rootworm adults, Fall armyworms, Flea beetles, Japanese beetles, Sap beetles, Southwestern corn borers, Leafhoppers	1¼ to 2½
	SPECIFIC DIRECTIONS: For optimum Chinch bug control, use ground equipment to apply at least 20 gals. of water per acre and direct spray toward stalk to provide thorough coverage.	
	European corn borers	1¾ to 2½
	SPECIFIC DIRECTIONS: For optimum European corn borer control, do not apply in less than 3 gals. of water per acre by air and 15 gals. of water by ground.	
Cutworms, Western bean cutworms		2½
	SPECIFIC DIRECTIONS: For optimum Cutworm control, apply in a 12-inch band over the row, using sufficient volume of water to obtain thorough coverage. For broadcast application, use at least 20 gals. by ground or 5 gals. by air per acre. For Cutworm control, this product is most effective against species which feed on the upper portions of the plant. For Western bean cutworms, treat when infestation averages 15% and at 90 to 100% tassel emergence. Treatment after 100% silk emergence will reduce effectiveness.	

**RESTRICTIONS AND PRECAUTIONS:** Observe Bee Caution. Do not apply more than a total of 10 lbs. per acre per crop. Do not apply within 48 days of harvest of grain and fodder or within 14 days of harvest or grazing of forage or silage.

#### GRAIN SORGHUM

Repeat applications as necessary up to a total of 4 times, but not more often than once every 7 days. Direct spray into forming heads for optimum control of insects attacking heads.

Crop	Pest	Lbs. of This Product Per Acre
Grain sorghum	Armyworms, Chinch bugs, Corn earworms, Fall armyworms, Stink bugs, Webworms	1¼ to 2½
	SPECIFIC DIRECTIONS: For optimum Chinch bug control, use high gallonage ground application at the base of the plants.	
	Southwestern corn borers	1¾
	Cutworms	2½
SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.		

**RESTRICTIONS AND PRECAUTIONS:** Do not apply within 21 days of harvest for grain or fodder or within 14 days of harvest or grazing of forage or silage. Do not apply more than a total of 7½ lbs. per acre per crop.

#### RICE

Crop	Pest	Lbs. of This Product Per Acre
Rice	Armyworms, Chinch bugs, Fall armyworms, Leafhoppers, Stink bugs	1¼ to 1¾
	SPECIFIC DIRECTIONS: Up to 2 applications per crop may be made, but not more often than once every 7 days.	
	Tadpole shrimp (CA Only)	1¾
SPECIFIC DIRECTIONS: For optimum Tadpole shrimp control, apply to water when pest first appears.		

**RESTRICTIONS AND PRECAUTIONS:** Do not apply within 14 days of harvest for grain or straw. Do not apply more than a total of 5 lbs. per acre per crop. May kill shrimp, crabs and crayfish. Do not apply propanil herbicides within 15 days before or after application of this product or plant injury may result.

#### SWEET CORN

Observe Bee Caution. Repeat applications as necessary up to a total of 8 times, but not more than once every 3 days. Optimum timing and good coverage are essential for effective control.

For insects attacking silks and ears, insecticide sprays should be applied starting when first silks appear and continuing until silks begin to dry. During silking, the minimum re-treatment interval (3 days) may not provide adequate levels of protection under conditions of rapid growth or severe pest pressure. The use of an alternative product should be considered in conjunction with this product.

Crop	Pest	Lbs. of This Product Per Acre
Sweet corn	Armyworms, Chinch bugs, Corn earworms, Corn rootworm adults, Fall armyworms, Flea beetles, Japanese beetles, Leafhoppers, Sap beetles, Southwestern corn borers	1¼ to 2½
	SPECIFIC DIRECTIONS: For optimum Chinch bug control, use ground equipment to apply at least 20 gals. of water per acre and direct spray toward stalk to provide thorough coverage.	
	European corn borers	1% to 2½
	SPECIFIC DIRECTIONS: For optimum European corn borer control, do not apply in less than 3 gals. of water per acre by air and 15 gals. of water by ground.	
	Cutworms, Western bean cutworms	2½
SPECIFIC DIRECTIONS: For optimum Cutworm control, apply in a 12-inch band over the row, using sufficient volume of water to obtain thorough coverage. For broadcast application, use at least 20 gals. by ground or 5 gals. by air per acre. For Cutworm control, this product is most effective against species which feed on the upper portions of the plant. For Western bean cutworms, treat when infestation averages 15% and at 90 to 100% tassel emergence. Treatment after 100% silk emergence will reduce effectiveness.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 2 days of harvest of ears, within 14 days of harvest or grazing of forage or within 48 days of harvest of fodder. Do not apply more than a total of 20 lbs. per acre per crop.		

#### WHEAT AND PROSO MILLET

Up to 2 applications per crop may be made, but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Wheat, Proso millet  (DO NOT USE IN CA)	Flea beetles	¾ to 1¼
	Cereal leaf beetles	1¼
	SPECIFIC DIRECTIONS: Application is effective against eggs, larvae and adults of the Cereal leaf beetles.	
	Armyworms, Fall armyworms	1¼ to 1¾
SPECIFIC DIRECTIONS: Application for Armyworm control should be made when Armyworms are actively feeding on the upper foliage and night temperatures are not expected to drop below 55°F. If applied by air to lush growth, use a minimum spray volume of 5 gals. per acre to optimize coverage.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 21 days of harvest for grain or straw or within 7 days of harvest or grazing for forage. Do not apply more than a total of 3¾ lbs. per acre per crop.		

#### CUCURBIT VEGETABLES

Repeat applications as necessary up to a total of 6 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Cucumbers, Melons, Pumpkins, Squash	Melonworms, Pickleworms	¾ to 1¼
	Cucumber beetles, Flea beetles, Leafhoppers, Squash bugs	1¼
	SPECIFIC DIRECTIONS: For optimum control of Squash bugs, apply sufficient spray volume for thorough coverage and time sprays for early morning or late afternoon.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 3 days of harvest. Do not apply more than a total of 7½ lbs. per acre per crop. Observe plant response precautions.		

#### FLAX

Up to 2 applications per crop may be made, but not more than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Flax (DO NOT USE IN CA)	Armyworms	1¼ to 1¾
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 42 days of harvest for Seed or Straw. Do not apply more than a total of 3¾ lbs. per acre per crop.		

#### FORAGE CROPS

On dense growth, use 25 to 40 gallons of water per acre with ground equipment to ensure adequate coverage.

Crop	Pest	Lbs. of This Product Per Acre
Alfalfa, Birdsfoot trefoil, Clovers	Blister beetles, Mexican bean beetles	¾ to 1¼
	Alfalfa caterpillars, Bean leaf beetles, Cucumber beetles, Green cloverworms, Japanese beetles, Leafhoppers, Potato leafhoppers, Threecornered alfalfa hoppers, Thrips, Velvetbean caterpillars	1¼
	Alfalfa blotch leafminers, Armyworms, Cloverhead weevils, Corn earworms, Cutworms, Egyptian alfalfa weevil larvae, Essex skipper, European alfalfa beetles, Fall armyworms, Lygus bugs, Stink bugs, Webworms, Yellowstriped armyworms	1¼ to 1¾
	SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
	Alfalfa weevil larvae (West of the Rocky Mountains)	1¼ to 1¾
	SPECIFIC DIRECTIONS: For Alfalfa weevil larvae, if pre-treatment damage is extensive, cut Alfalfa and treat the stubble. This product is not effective against adult Alfalfa weevils.	
	Alfalfa weevil larvae (East of the Rocky Mountains)	1¾
SPECIFIC DIRECTIONS: For Alfalfa weevil larvae, if pre-treatment damage is extensive, cut Alfalfa and treat the stubble. This product is not effective against adult Alfalfa weevils.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Observe Bee Caution. Observe plant response precautions. Do not apply more than once per cutting. Do not apply within 7 days of harvest or grazing. Do not exceed 1¾ lbs. per acre per cutting. Carbaryl may cause a temporary bleaching of tender Alfalfa foliage.		

#### PASTURE AND GRASSES GROWN FOR SEED

Up to 2 applications per year may be made, but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Pasture and Grasses grown for seed	Armyworms, Chinch bugs, Essex skippers, Fall armyworms, Striped grass loopers, Thrips, Range caterpillars, Range crane flies, Ticks	1¼ to 1¾
SPECIFIC DIRECTIONS: To control Thrips in grasses grown for seed, use high spray pressure to improve penetration into the boot. Carefully mark swaths to avoid over-application.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of harvest or grazing. Do not exceed a total of 3¾ lbs. per acre per year.		

#### RANGELAND

Do not make more than 1 application per year. Carefully mark swaths to avoid over-application.

Crop	Pest	Lbs. of This Product Per Acre
Rangeland	Black grass bugs, Grasshoppers, Mormon crickets, Range caterpillars, Range crane flies	¾ to 1¼
	Ticks	1¼
<b>RESTRICTIONS AND PRECAUTIONS:</b> May be harvested or grazed the same day of treatment. Do not apply more than 1¼ lbs. per acre per year.		

**FRUITING VEGETABLES**

Repeat applications as necessary up to a total of 7 times, but not more often than once every 7 days. When disease transmission is suspected, monitor fields following application and re-treat if reinfestation occurs, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Eggplant, Peppers, Tomatoes	Colorado potato beetles, European corn borers, Fall armyworms, Lace bugs, Stink bugs (suppression), Tarnished plant bugs, Thrips (suppression), Tomato fruitworms, Tomato hornworms, Tomato pinworms	1¼ to 2½
	SPECIFIC DIRECTIONS: Thorough coverage is essential to effectively suppress Stink bugs.	
	Flea beetles, Leafhoppers	¾ to 1¼
	Cutworms	2½
SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 3 days of harvest. Do not apply more than a total of 10 lbs. per crop.		

**LEAFY VEGETABLES**

Repeat applications as necessary up to a total of 5 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Celery, Dandelion, Endive, Lettuce (Head and Leaf), Parsley, Spinach, Swiss chard	Flea beetles, Harlequin bugs, Leafhoppers	¾ to 1¼
	Armyworms, Aster leafhoppers, Corn earworms, Fall armyworms, Imported cabbageworms, Lygus bugs, Spittlebugs, Stink bugs, Tarnished plant bugs	1¼ to 2½
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of harvest. Do not apply more than a total of 7½ lbs. per acre per crop.		

**LEGUME VEGETABLES**

Repeat applications as necessary up to a total of 4 times, but not more often than once every 7 days. Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects and spiders. Use the higher rates for heavy populations and larger instars.

Crop	Pest	Lbs. of This Product Per Acre
Cowpeas, Fresh and Dried beans ( <i>Phaseolus</i> species including Kidney, Navy and Snap), Fresh and Dried peas ( <i>Pisum</i> species), Lentils, Southern peas, Soybeans	Bean leaf beetles, Blister beetles, Cucumber beetles, Grape colaspis, Green cloverworms, Japanese beetles, Mexican bean beetles, Velvetbean caterpillars	¾ to 1¼
	Corn earworms	¾ to 1½
	Alfalfa caterpillars, Colorado potato beetles, Flea beetles, Leafhoppers, Threecornered alfalfa hoppers, Thrips, Western bean cutworms	1¼
(DO NOT USE ON LENTILS IN CA)	Armyworms, Cutworms, European corn borers, Fall armyworms, Stink bugs, Tarnished plant bugs, Webworms	1¼ to 1½
SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.		
	Alfalfa loopers (suppression), Cowpea curculios (suppression), Painted ladies (Thistle caterpillars), Pea leaf weevils, Pea weevils, Saltmarsh caterpillars, Woollybear caterpillars, Yellowstriped armyworms	1½
	CA ONLY: Corn earworms (suppression), Limabean podborers (suppression), Lygus bugs (suppression), Stink bugs (suppression)	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of Fresh beans or peas or within 21 days of harvest of Dried beans or peas, seed or hay. Do not apply more than a total of 7½ lbs. per acre per crop. Do not apply a combination of this product and 2,4-DB herbicides to Soybeans as crop injury may result. Observe plant response precautions.		

**NON-CROPLAND**

Up to 2 applications per year may be made, but not more often than once every 14 days. Carefully mark swaths to avoid over-application.

Site	Pest	Lbs. of This Product Per Acre
Conservation Reserve Program Acreage, Set-Aside Program Acreage, Wasteland, Right-of-Ways, Hedge-rows, Ditchbanks, Roadsides	Black grass bugs	½ to ¾
	Mormon crickets, Range caterpillars, Range crane flies	¾ to 1¼
	Ticks	1¼ to 1½
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of grazing or harvest for forage or hay. Do not apply more than a total of 3¾ lbs. per acre per year.		

**OKRA**

Apply using ground equipment in sufficient volume to ensure good coverage. For Grasshopper control, refer to "CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES" section.

Crop	Pest	Lbs. of This Product Per Acre
Okra	Corn earworms, Stink bugs	1¼ to 1½
	SPECIFIC DIRECTIONS: Treat on a 6 to 8-day interval as needed.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 3 days of harvest. Do not apply more than a total of 7½ lbs. per acre per season.		

**PEANUTS**

Repeat applications as necessary up to a total of 5 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Peanuts	Blister beetles, Mexican bean beetles	¾ to 1¼
	Alfalfa caterpillars, Bean leaf beetles, Cucumber beetles, Green cloverworms, Japanese beetles, Leafhoppers, Rednecked peanutworms, Threecornered alfalfa hoppers, Thrips, Velvetbean caterpillars	1¼
	SPECIFIC DIRECTIONS: For optimum control of Thrips, use directed or banded sprays with hollow cone spray nozzles. Ensure adequate coverage for the underside of leaves.	
	Armyworms, Corn earworms, Fall armyworms, Stink bugs, Webworms	1¼ to 1½
	Whitefringed beetle adults, Cutworms	2½
SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Observe plant response precautions. Do not apply within 14 days of harvest. Do not apply more than a total of 10 lbs. per acre per crop.		

**PRICKLY PEAR CACTUS**

Apply using ground equipment in sufficient volume to ensure good coverage. For Grasshopper control, refer to "CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES" section.

Site	Pest	Lbs. of This Product Per Acre
Prickly pear cactus	Cochineal scales (crawlers)	2½
	SPECIFIC DIRECTIONS: Apply as needed at 7- to 10-day intervals.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 3 days of harvest. Do not apply more than 7½ lbs. per acre per season.		

**ROOT AND TUBER CROPS**

(Except Sugar beets and Sweet potatoes)  
Repeat applications as necessary up to a total of 6 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Carrots, Garden beets, Horseradish, Parsnips, Potatoes, Radishes, Rutabagas, Salsify	Flea beetles, Leafhoppers	¾ to 1¼
	Armyworms, Aster leafhoppers, Colorado potato beetles, Corn earworms, Cutworms, European corn borers, Fall armyworms, Lace bugs, Lygus bugs, Spittlebugs, Stink bugs, Tarnished plant bugs	1¼ to 2½
	SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 7 days of harvest. Do not apply more than a total of 7½ lbs. per acre per crop.		

### SUGAR BEETS

Repeat applications as necessary up to a total of 2 times, but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Sugar beets	Armyworms, Beet leaf beetles, Fall armyworms, Flea beetles, Leafhoppers, Webworms	1¼ to 1½
	Cutworms	1½
	SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 28 days of harvest for roots or forage. Do not apply more than a total of 5 lbs. per acre per crop.		

### SWEET POTATOES

For foliar sprays, repeat applications as necessary up to a total of 8 times, but no more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Sweet potatoes	Corn earworms, Cucumber beetles, Flea beetles, Sweet potato hornworms, Sweet potato weevils, Tortoise beetles, Whitefringed beetles	1¼ to 2½
	SPECIFIC DIRECTIONS: Pre-plant dip for control of Sweet potato weevils: Just prior to planting, dip Sweet potato cuttings in a suspension containing 10 lbs. of this product in 100 gals. of water (1.6 ozs. per gal. of water).	
	Yellowstriped armyworms	2½
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 7 days of harvest. Do not apply more than a total of 10 lbs. per acre per crop with in-season sprays. Do not apply more than a total of 1½ lbs. per acre as a pre-plant dip treatment.		

### SMALL FRUITS AND BERRIES

Repeat applications as necessary up to a total of 5 times, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Blueberries, Caneberries, Cranberries, Grapes, Strawberries	European fruit lecaniums, European raspberry aphids, Flea beetles, Grape leafhoppers, Grape leafrollers, Japanese beetles, Leafhoppers, Leafrollers, Meadow spittlebugs, Omnivorous leafhoppers, Rose chafers, Snowy tree crickets, Strawberry bud weevils, Strawberry clippers, Strawberry fruitworms, Strawberry leafrollers, Strawberry weevils, Western grapeleaf skeletonizers, Western yellowstriped armyworms	1¼ to 2½
	SPECIFIC DIRECTIONS: In Grapes for Grape leafroller control, apply before the first brood larvae emerge from rolls.	
	Blueberry maggots, Cherry fruitworms, Cranberry fireworms, Cranberry fruitworms, Cranberry twig girdlers, Elm spanworms, Gypsy moths, Sparganothis worms, Tarnished plant bugs	1½ to 2½
	Eightspotted foresters, Cutworms, Grape berry moths, June beetles, Omnivorous leafrollers, Orange tortrix, Raspberry fruitworms, Raspberry sawflies, Redbanded leafrollers, Saltmarsh caterpillars	2½
	SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Observe Bee Caution. Do not apply within 7 days of harvest. Do not apply more than a total of 12½ lbs. per acre per crop. <b>Note:</b> Use in Cranberries may kill shrimp and crabs. Do not use in areas where these are important resources. Carbaryl may injure Early dawn and Sunrise varieties of Strawberries. In Grapes, do not concentrate spray on the bunch or visible residues may result.		

### SUNFLOWERS

Up to 2 applications may be made, but not more often than once every 7 days.

Crop	Pest	Lbs. of This Product Per Acre
Sunflowers  (DO NOT USE IN CA)	Stem weevils, Sunflower beetles	1¼ to 1½
	Armyworms, Cutworms, Fall armyworms, Sunflower moths	1½
	SPECIFIC DIRECTIONS: For Cutworm control, this product is most effective against species which feed on the upper portions of the plant.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 30 days of grazing or harvest for forage or within 60 days of harvest for seed. Do not apply more than a total of 3½ lbs. per acre per crop.		

### TOBACCO

**Plant bed and Field Treatment**—Repeat treatments as necessary up to a total of 4 times per crop, but not more often than once every 7 days. Use lower rate on young plants (up to knee height). Use at least 10 gallons of prepared spray per acre. Begin treatments when worms are small.

Crop	Pest	Lbs. of This Product Per Acre
Tobacco	Budworms, Fall armyworms, Hornworms, Japanese beetles, June beetles, Suckflies, Tobacco flea beetles	1¼ to 2½
<b>RESTRICTIONS AND PRECAUTIONS:</b> Tobacco may be harvested on the day of treatment. Do not apply more than a total of 10 lbs. per acre per crop. Observe plant response precautions.		

### TREE FRUIT CROPS

On all tree fruit crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, density and other factors. Repeat applications as necessary up to a total of 8 times, but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Citrus fruits	Avocado leafrollers, California orangedogs, Citrus cutworms, Fruitree leafrollers, Orange tortrix, Western tussock moths	2½ to 3½
	Citrus rust mites, Eriophyid mites, Plant bugs, Scale insects (Black scales, Brown soft scales, California red scales [except in CA], Citrus snow scales, Yellow scales [except in CA])	3¾ to 6¼
	SPECIFIC DIRECTIONS: For Scale control, apply when crawlers are present. For best control of Eriophyid mites, including Citrus rust mites, apply when pest populations are low.	
	Apopka weevils (adult), Citrus root weevils (adults), Fuller rose beetles, Little leaf notchers (adult)	6¼ to 9¾
	CA Only: California red scales, Yellow scales	6¼ to 20
SPECIFIC DIRECTIONS: Do not make more than 1 application per season for California red scales. Apply when crawlers are present.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Observe Bee Caution. Do not apply within 5 days of harvest. Do not apply more than a total of 25 lbs. per acre per crop.		

### OLIVES

Up to 2 applications per crop may be made, but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Olives	Scale insects (Black scales, Olive scales)	6¼ to 9¾
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of harvest. Do not apply more than a total of 18½ lbs. per acre per crop.		

**POME FRUITS**

Repeat applications as necessary up to a total of 8 times per crop (including thinning sprays on Apples), but not more often than once every 14 days.

Crop	Pest	Lbs. of This Product Per Acre
Apples, Crabapples, Pears, Loquats, Oriental pears	White apple leafhoppers	5/8 to 1 1/8
	Apple aphids, Codling moths	1 1/4 to 3 3/4
	Apple aphids, Apple maggots, Apple mealybugs, Apple rust mites, Bagworms, California pearslugs (Pear sawflies), European apple sawflies, Eyespotted bud moths, Fruittree leafrollers, Gypsy moths, Japanese beetles, Lesser appleworms, Lygus bugs, Orange tortrix, Pearleaf blister mites, Pear psyllas, Pear rust mites, Periodical cicadas, Plum curculios, Redbanded leafrollers, Rosy apple aphids, Scale insects (Forbes scales, Lecanium scales, San Jose scales), Tarnished plant bugs, Tentiform leafminers, Woolly apple aphids, Yellowheaded fireworms	1 1/8 to 3 3/4
SPECIFIC DIRECTIONS: On Apples, avoid use during the period from full bloom until 30 days after full bloom unless fruit thinning is desired. Use for pest control during this period also may result in fruit removal. For Psylla control, apply when eggs hatch or young nymphs are present. For Scale control, apply when crawlers are present.		
Apples Only, for fruit thinning	--	1 1/4 to 3 3/4
SPECIFIC DIRECTIONS: Observe Bee Caution. For easily thinned varieties: Apply 1/8 to 5/8 lbs. per 100 gals. of spray mixture. For difficult-to-thin varieties: Apply 5/8 to 1 1/4 lbs. per 100 gals. of spray mixture. Apply between 10 and 25 days after full bloom. Factors such as tree age, variety, nutrition, previous crop, pruning, bloom and degree of set favor excessive fruit thinning with this product. Exercise caution to avoid possible yield reduction. Rates may vary depending on variety and local orchard conditions. Consult with your county extension service or other experts for advice on the proper use of this product. PRECAUTIONS: The use of this product may result in fruit deformity under certain environmental conditions. Before using on any variety of Apples, the user must weigh the risk versus benefits when using this product, particularly when using between 80% petal fall and 6 mm fruit size. Red delicious are more sensitive to this phenomenon and in particular, the varieties Bisbee, Red chief and Vallee spur are very susceptible to conditions causing fruit deformity. Precipitation and temperatures below 65°F increases the possibility of fruit deformity. The use with Summer spray oils and wetting agents may increase the risk of fruit deformity and injury.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply to Quince. Do not use on Pears between the tight flower cluster up to the 20 mm fruit size. Use during this period may result in undesirable fruit thinning and/or deformed fruit. Do not apply within 3 days of harvest. Do not apply more than a total of 18 3/4 lbs. per acre per crop. Do not make more than a total of 8 applications per crop. <b>FOR PROTECTION OF HONEYBEES:</b> Remove all bee hives from orchard to be treated prior to application. Do not apply this product if bees are actively foraging in orchard. If weed bloom is present, mow the cover crop on the orchard floor prior to applying this product.		

**STONE FRUITS**

Crop	Pest	Lbs. of This Product Per Acre
Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Apple pandemis, Black cherry aphids, Cherry fruitworms, Cherry maggots (Cherry fruit flies), Codling moths, Cucumber beetles, Eastern tent caterpillars, Eyespotted bud moths, European earwigs, Fruittree leafrollers, Green fruitworms, Gypsy moths, Japanese beetles, June beetles, Lesser peachtree borers, Mealy plum aphids, Orange tortrix, Oriental fruit moths, Peach twig borers, Periodical cicadas, Plum curculios, Prune leafhoppers, Redbanded leafrollers, Rose chafers, Scale insects (Brown soft scales, Forbes scales, Lecanium scales, Olive scales, Oystershell scales, San Jose scales), Tarnished plant bugs, Tussock moths, Variegated leafrollers	2 1/2 to 3 3/4
	SPECIFIC DIRECTIONS: Observe Bee Caution. Repeat applications as necessary up to a total of 3 times per crop, but not more often than once every 7 days. An additional application at the dormant or delaye- dormant timing may be made. For optimum Scale control, apply when crawlers are present. For Lesser peachtree borers, best results have been found by thoroughly spraying limbs and tree trunks at weekly intervals during moth flight.	
	CA ONLY: Black cherry aphids, Cherry fruitworms, Cherry maggots (Cherry fruit flies), Codling moths, Cucumber beetles, Eyespotted bud moths, European earwigs, Fruittree leafrollers, Green fruitworms, Mealy plum aphids, Orange tortrix, Oriental fruit moths, Peach twig borers, Scale insects (Brown soft scales, Forbes scales, Lecanium scales, Olive scales, Oystershell scales, San Jose scales), Tarnished plant bugs, Tussock moths	3 3/4 to 5
	SPECIFIC DIRECTIONS: Observe Bee Caution. Repeat applications as necessary up to a total of 3 times per crop but not more often than once every 14 days. An additional application at the dormant or delaye- dormant timing may be made. For optimum Scale control, apply when crawlers are present.	
	Peach twig borers, Scale insects (Brown soft scales, Forbes scales, Lecanium scales, Olive scales, Oystershell scales, San Jose scales)	5 to 6 1/4
SPECIFIC DIRECTIONS: For dormant or delayed-dormant timing, apply in combination with a recommended dormant oil. Refer to the dormant oil product label for additional use directions and restrictions.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 3 days of harvest, except in CA. In CA, do not apply within 1 day of harvest. Do not apply more than a total of 17 1/2 lbs. per acre per crop. Do not apply more than a total of 6 1/4 lbs. per acre at the dormant or delayed-dormant timing. Do not apply more than a total of 11 1/4 lbs. per acre during the production season.		

**TREE NUT CROPS**

On all tree nut crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, density and other factors.

Crop	Pest	Lbs. of This Product Per Acre
Pistachios	Brown soft scales, Lecanium scales, Navel orangeworms	3¾ to 6¼
	SPECIFIC DIRECTIONS: Repeat applications as necessary up to a total of 4 times per crop (including any applications at the dormant or delayed-dormant timing), but not more often than once every 7 days. For Scale control, apply when crawlers are present.	
	Scale insects	5 to 6¼
	SPECIFIC DIRECTIONS: For dormant or delayed-dormant timing, apply in combination with a recommended dormant oil. Refer to the dormant oil product label for additional use directions and restrictions.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not apply within 14 days of harvest. Do not apply more than a total of 18¾ lbs. per acre per crop, including any application at the dormant or delayed-dormant timing.		
Almonds, Chestnuts, Filberts, Pecans, Walnuts	Black margined aphids, Calico scales, Codling moths, European fruit lecaniums, Fall webworms, Filbert aphids, Filbert leafrollers, Filbertworms, Frosted scales, Fruittree leafrollers, Hickory shuckworms, Lesser webworms, Navel orangeworms, Peach twig borers, Pecan leaf phylloxeras, Pecan stem phylloxeras, Pecan nut casebearers, Pecan spittlebugs, Pecan weevils, San Jose scales, Twig girdlers, Walnut caterpillars	2½ to 6¼
	SPECIFIC DIRECTIONS: Observe Bee Caution. Repeat applications as necessary up to a total of 4 times per crop (including any applications at the dormant or delayed-dormant timing), but not more often than once every 7 days. Use lower rates for pests attacking leaves. Use higher rates for pests attacking fruit and for higher infestations. For Scale control, apply when crawlers are present. For Peach twig borers, best results with foliar applications have been found by making applications in popcorn or petal fall stages when the May brood begins to hatch. For Navel orangeworms in Almonds and Walnuts, best results have been found by timing early and mid-season applications to correspond with moth flight peaks. For Filbert leafrollers, best results have been found by making applications when eggs are hatching, repeating application in first appearance of moths and again 3 to 4 weeks later. For Codling moths in Walnuts, best results have been found by making applications when average cross-sectional diameters of developing nuts are ½ to ¾ inch and again during middle or late June as needed.	
	Chestnut weevils, European earwigs	5 to 6¼
	SPECIFIC DIRECTIONS: For Chestnut weevils, best results have been found with 4 applications at weekly intervals beginning in late July. The last application should be made prior to shuck split. For European earwigs, thorough coverage of trunks, branches and nuts is needed for best results.	
	RESTRICTIONS AND PRECAUTIONS: Do not apply within 14 days of harvest. Do not apply more than a total of 18¾ lbs. per acre per crop, including any applications at the dormant or delayed-dormant timing.	
Almonds Only	Peach twig borers, Scale insects	5 to 6¼
	SPECIFIC DIRECTIONS: For dormant or delayed-dormant timing, apply in combination with a recommended dormant oil. Refer to the dormant oil product label for additional use directions and restrictions.	

**FORESTED AREAS AND RANGELAND TREES**

Apply in sufficient volume for adequate coverage. This will vary depending on the tree size, density and stage of growth.

Crop	Pest	Lbs. of This Product Per Acre
Non-urban forests, Tree plantations, Planted Christmas trees, Parks, Rural shelter belts, Rangeland trees	Ants, Apple aphids, Armyworms, Ash whiteflies, Azalea leafminers, Bagworms, Balsam twig aphids, Birch leafminers, Blister beetles, Boxelder bugs, Boxwood leafminers, Browntail moths, Cankerworms, Catalpa sphinx, Chiggers, Cooley spruce gall adelgids, Cutworms, Cypress tip moths, Douglas-fir tussock moths, Eastern spruce gall adelgids, Elm leaf aphids, Elm leaf beetles, Elm spanworms, Eriophyid mites, European pine shoot moths, Fall armyworms, Flea beetles, Fuchsia gall mites, Fuller rose beetles, Gall midges, Gall wasps, Greenstriped mapleworms, Grasshoppers, Hackberry nipplegall makers, Holly bud moths, Holly leafminers, Jackpine budworms, Japanese beetles, Jeffrey pine needleminers, June beetles, Lace bugs, Leafhoppers, Leafrollers, Locust borers, Maple leafcutters, Mealybugs, Mimosa webworms, Nantucket pine tip moths, Oak leafminers, Oak moths, Oak skeletonizers, Oakworm complex, Oleander caterpillars, Olive ash borers, Orangestriped oakworms, Periodical cicadas, Pine loopers, Pine sawflies, Pine spittlebugs, Pitch pine tip moths, Plant bugs, Poinsettia hornworms, Psyllids, Puss caterpillars, Redhumped oakworms, Rose aphids, Rose chafers, Rose slugs, Saddled prominents, Sawflies (exposed), Scale insects (crawlers), Sowbugs, Spiney elm caterpillars, Springtails, Spruce budworms, Spruce needleminers, Subtropical pine tip moths, Tent caterpillars, Thornbugs, Thrips (exposed), Ticks, Walnut caterpillars, Webworms, Western hemlock loopers, Western spruce budworms, Willow leaf beetles, Woolly gall aphids, Yellow-poplar weevils	1¼
	SPECIFIC DIRECTIONS: Observe plant response precautions. Obtain thorough coverage of upper and lower leaf surfaces. The addition of a sticker may improve residual control. To control Scale insects, treat trunks, stems and twigs in addition to plant foliage. For optimum Worm control, treat when pests are small. Do not use on syrup-producing Sugar maples where sap is harvested. Applications for control of Maple leafcutters on Sugar maple should be made when larvae are in 2nd instar after mining and as cases are being formed. Repeat treatments as necessary up to a total of 2 times per year, but not more often than once every 7 days.	
	Gypsy moths	¾ to 1¼
	SPECIFIC DIRECTIONS: For Gypsy moth control, use the higher rate for heavy infestations.	
Elm bark beetles, Ips engraver beetles, Mountain pine beetles, Roundheaded pine beetles, Spruce beetles, Western pine beetles	2% solution (3 ozs. per gal.)	
	SPECIFIC DIRECTIONS: Direct Trunk Treatment - Effective as a preventative treatment only. Repeat annually as required to prevent Beetle attacks. Apply 1 gal. of spray per 50 sq. ft. of bark prior to Beetle flight or host-tree attack. Treat tree trunk from ground level up, until trunk diameter is less than 5 inches. For Elm bark beetles; apply approximately 20 to 30 gals. of spray mixture for each 50 ft. of Elm tree for thorough coverage of all bark surfaces on trunks, limbs and twigs. Do not make more than 2 applications per year or repeat applications more often than once every 6 months.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not make more than 2 applications per year. To prepare small volumes of spray mixture, use 6.5 grams (approx. 1.2 tbsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1¼ lbs. per acre are indicated. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.		



**Note:** All references to Armyworms on the plants listed below refer to the species *Pseudaletia unipuncta*, often called the "True armyworm". Except where indicated otherwise, this product is not registered for control of other Armyworm species. Regional differences have been noted in the susceptibility of certain strains of Fall armyworms to carbaryl. If local experience indicates inadequate control, use an alternative pesticide.

**TREES AND ORNAMENTALS**

Apply in sufficient volume for adequate coverage. This will vary depending on the plant or tree size, density and stage of growth. Do not use on Boston ivy, Virginia creeper and Maidenhair fern. During early season it may also injure Virginia and Sand pines.

Crop	Pest	Lbs. of This Product Per Acre
Trees, Ornamentals: (Roses, Flowers, Other Herbaceous plants), Woody plants, Shrubs in commercial and residential areas and in Nursery production areas	Ants, Apple aphids, Armyworms, Ash whiteflies, Azalea leafminers, Bagworms, Balsam twig aphids, Birch leafminers, Blister beetles, Boxelder bugs, Boxwood leafminers, Browntail moths, Cankerworms, Catalpa sphinx, Chiggers, Cooley spruce gall adelgids, Cutworms, Cypress tip moths, Douglas-fir tussock moths, Eastern spruce gall adelgids, Elm leaf aphids, Elm leaf beetles, Elm spanworms, Eriophyid mites, European pine shoot moths, Fall armyworms, Flea beetles, Fuchsia gall mites, Fuller rose beetles, Gall midges, Gall wasps, Greenstriped mapleworms, Grasshoppers, Hackberry nipplegall makers, Holly bud moths, Holly leafminers, Jackpine budworms, Japanese beetles, Jeffrey pine needleminers, June beetles, Lace bugs, Leafhoppers, Leafrollers, Locust borers, Maple leafcutters, Mealybugs, Mimosa webworms, Nantucket pine tip moths, Oak leafminers, Oak moths, Oak skeletonizers, Oakworm complex, Oleander caterpillars, Olive ash borers, Orange-striped oakworms, Periodical cicadas, Pine loopers, Pine sawflies, Pine spittlebugs, Pitch pine tip moths, Plant bugs, Poinsettia hornworms, Psyllids, Puss caterpillars, Redhumped oakworms, Rose aphids, Rose chafers, Rose slugs, Saddled prominents, Sawflies (exposed), Scale insects (crawlers), Sowbugs, Spiney elm caterpillars, Springtails, Spruce budworms, Spruce needleminers, Subtropical pine tip moths, Tent caterpillars, Thornbugs, Thrips (exposed), Ticks, Walnut caterpillars, Webworms, Western hemlock loopers, Western spruce budworms, Willow leaf beetles, Woolly gall aphids, Yellow-poplar weevils	1¼
SPECIFIC DIRECTIONS: Observe plant response precautions. Obtain thorough coverage of upper and lower leaf surfaces. The addition of a sticker may improve residual control. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. For optimum Worm control, treat when pests are small. Do not use on syrup-producing Sugar maples where sap is harvested. Applications for control of Maple leafcutters on Sugar maple should be made when larvae are in 2nd instar after mining and as cases are being formed. Repeat treatments as necessary up to a total of 2 times per year, but not more often than once every 7 days.		
Gypsy moths		9/10 to 1¼
SPECIFIC DIRECTIONS: For Gypsy moth control, use the higher rate for heavy infestations.		
Elm bark beetles, Ips engraver beetles, Mountain pine beetles, Roundheaded pine beetles, Spruce beetles, Western pine beetles		2% solution (3 ozs. per gal.)
SPECIFIC DIRECTIONS: Direct Trunk Treatment - Effective as a preventative treatment only. Repeat annually as required to prevent Beetle attacks. Apply 1 gal. of spray per 50 sq. ft. of bark prior to Beetle flight or host-tree attack. Treat tree trunk from ground level up, until trunk diameter is less than 5 inches. For Elm bark beetle; apply approximately 20 to 30 gals. of spray mixture for each 50 ft. of Elm tree for thorough coverage of all bark surfaces on trunks, limbs and twigs. Do not make more than 2 applications per year or repeat applications more often than once every 6 months.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not make more than a total of 6 foliar and 2 direct trunk treatments per year. To prepare small volumes of spray mixture, use 6.5 grams (approx. 1.2 tbsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1¼ lbs. per acre are indicated. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.		

**TURFGRASS**

Apply the recommended rates using pressure-type equipment in sufficient spray volume for thorough coverage and thatch penetration.

Crop	Pest	Lbs. of This Product Per Acre
Golf turf, Sports fields, Sod farms, Domestic and Commercial lawns, Cemeteries, Parks, Campsites, Recreational areas	Ants, Armyworms, Centipedes, Chiggers, Cutworms, Earwigs, Essex skippers, European chafers, Fall armyworms, Fiery skippers, Grasshoppers, Green June beetle larvae, Leafhoppers, Lucerne moths, Millipedes, Mosquitoes (adults), Scarab beetle adults (May beetles, June beetles, Japanese beetles, Green June beetles), Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped armyworms	2½ to 5
SPECIFIC DIRECTIONS: For Armyworm, Cutworm and Fall armyworm control: Do not irrigate treated areas within 24 hours following insecticide application. For Green June beetle grub (larvae) control: Make applications when Grubs are feeding near the soil surface. Water or irrigate Turfgrass soon after treatment. Repeat applications as necessary up to a total of 4 times a year, but not more often than once every 7 days.		
Chinch bugs, Sod webworms (Lawn moths)		7½ to 10
SPECIFIC DIRECTIONS: For Chinch bug control: Treat entire grass area rather than just damaged areas. Irrigation prior to treatment will aid in penetration of insecticide into Turfgrass. Do not irrigate treated areas within 24 hours following insecticide application. Repeat applications as necessary up to 2 times per year, but not more often than once every 7 days.		
Bluegrass billbug larvae, European crane flies, Fleas, White grubs (such as Japanese beetles, Chafer beetles and <i>Phyllophaga</i> spp. larvae), Black turfgrass ataenius beetle larvae, Hyperodes weevils (Bluegrass weevil) larvae		10
SPECIFIC DIRECTIONS: For European crane fly control: Treatments should be applied in early Spring (April 1 to April 15) or as recommended by local Agricultural Extension Service Agents. Water or irrigate Turfgrass soon after treatment. For White grub control: Applications should be made when Grubs are feeding near the soil surface, usually during late March through May or July through early September or as recommended by local Agricultural Extension Service Agent. Water or irrigate Turfgrass soon after treatment. Repeat applications as necessary up to 2 times per year, but not more often than once every 7 days.		
<b>RESTRICTIONS AND PRECAUTIONS:</b> DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.		

**CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES**

Crop	Pest	Lbs. of This Product Per Acre
All crops on this label	Grasshoppers	5% to 17%*
SPECIFIC DIRECTIONS: Apply 5% to 9/10 lb. per acre of this product for nymphs on small plants or sparse vegetation. Apply 1¼ to 17% lbs. per acre for mature Grasshoppers or applications to dense foliage or if extended residual control is desired. Be certain spray volumes are appropriate to assure adequate coverage.		
<b>*Note:</b> Refer to individual site listing elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions.		

**CONTROL OF TICKS WHICH VECTOR LYME DISEASE**

For control of juvenile and adult Ticks which vector Lyme disease, apply the recommended amount in sufficient volume for thorough coverage.

Crop	Pest	Lbs. of This Product Per Acre
All crops on this label: Pastures, Forested areas, Wasteland, Right-of-Ways, Hedges, Ditchbanks, Roadsides, Set-aside and Conservation Reserve Program acreage, Trees and Ornamentals, Turfgrass, Lawns and Recreational Turfgrasses (including Lawns and Perimeters, Golf courses, Sports fields, Cemeteries, Parks and Pastures, Shrubs, Wooded areas, (including: Military posts, Logging camps and Campsites)	<i>Ixodes</i> spp. (Deer ticks, Bear ticks, Black legged ticks), <i>Amblyomma</i> spp. (Lone star ticks)	1¼ to 2½*
	SPECIFIC DIRECTIONS: Use the higher rate for heavy Tick infestations*. Use higher spray volumes for dense ground cover or heavy leaf litter. Target applications for nymphal control in late Spring or early Summer. Control of adult Ticks can be obtained with late Summer and Fall applications. Do not use spot treatments. Treat entire area and perimeter areas where exposure to Ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Re-treat as necessary to maintain adequate control levels**.	

**RESTRICTIONS AND PRECAUTIONS**

\*\*Note: For control of Ticks which vector Lyme disease: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED.

**IMPORTED FIRE ANTS**

Crop/Site	Pest	Lbs. of This Product Per Volume of Water
Pastures, Rangelands, Forested areas, Trees and Ornamentals, Turfgrass, Wastelands, Lawns, Cemeteries, Recreational areas (including Turf, Golf courses and Parks)	Imported fire ants	8.4 grams per gal.
	SPECIFIC DIRECTIONS: Drench Application - Apply a total of 2 gals. of the diluted solution over the surface of each mound or at least 1 qt. per 6 inches of mound diameter using a bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding areas to a 4-ft. diameter (12 sq. ft.). Do not disturb mound prior to treatment. Pour solution from a height of about 3 ft. to give sufficient force to break mound apex and flow into Ant tunnels. For best results apply in cool weather (65 to 80°F) or in early morning or late evening hours. Repeat application if mound activity resumes after 7 days. Treat new mounds as they appear. Pressurized sprays may disturb the Ants and cause migration, reducing product effectiveness.	
Nursery stock, Vegetable transplants*, Foliage plants, Bedding plants (Outdoor use only)	Imported fire ants	1½ lbs. per 100 gals.
	SPECIFIC DIRECTIONS: Avoid contact with foliage and treat only the growing media when using on bedding plants. Do not make more than one application, either as a root dip or drench treatment (applied to the point of saturation).	

**RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL DRENCH HAS DRIED. DO NOT USE IN GREENHOUSES. To prepare small amounts, use 8.4 grams (approx. 1% tbsps.) of this product per each gal. of mix where 1½ lbs. per 100 gals. are indicated. \*Note: DO NOT USE ON ANY FOOD CROP NOT LISTED ON THIS LABEL. Refer to specific crop section for additional restrictions and precautions.

**ADULT MOSQUITO CONTROL**

Apply in sufficient gallonage for thorough coverage.

Crop	Pest	Lbs. of This Product Per Acre
Pastures, Rangeland, Forested lands, Trees and Ornamentals, Turfgrass, Wastelands	Mosquitoes (adults)	½ to 1¼*
SPECIFIC DIRECTIONS: Observe Bee Caution. Treat shrubbery and areas where adult Mosquitoes congregate. Treat when adult Mosquitoes are active in early mornings or late evenings. Repeat applications as necessary*. Use ½ to ¾ lb. per 100 gals. in mistblowers, 1¼ lbs. per acre in low pressure ground sprayers, or ¾ to 1¼ lbs. per acre in aerial sprays.		

**RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED. May kill shrimp and crabs. Do not use in areas where these are important resources. To prepare small volumes of spray mixture, use 6.5 grams (approx. 1.2 tbsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1¼ lbs. per acre are indicated. Note: Refer to individual site listing elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions.

**NUISANCE PEST CONTROL**

For use as a perimeter treatment to prevent infestations of buildings, apply the specified dosage in a band 6 to 10 feet wide around the outside perimeter of the structure.

Pest	Rate of This Product
Ants, Bees, Brown dog ticks, Centipedes, Cockroaches, Crickets, Darkling beetles (Lesser mealworms or Litter beetles), Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Spiders, Wasps	1% solution or 1½ ozs. per gal.

SPECIFIC DIRECTIONS: Observe plant response precautions. Apply as a coarse, wet spray in a band 6 to 10 ft. wide around the outside perimeter of buildings. Confine applications to areas immediately adjacent to the building. Direct application to the structure should be minimal and restricted to cracks, crevices and areas where insects tend to congregate. For use in and around flower beds and ornamental plantings, apply as a coarse, wet spray in and around plant beds. May be applied to outdoor sleeping quarters of pets. Do not treat animals. Do not allow animals to use treated area until sprays have dried.

**RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED. Note: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Spray deposits on painted or stained surfaces of finishes (e.g., cars, house trailers, boats, etc.) should be removed immediately by washing to prevent discoloration. Avoid application to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

**POULTRY INSECT CONTROL**

Site	Pest	Lbs. of This Product Per 100 Gals. of Water
Direct spray on Chickens, Ducks, Geese, Game Birds, Pigeons, Turkeys	Chicken mites, Fleas, Lice, Northern fowl mites	--
SPECIFIC DIRECTIONS: For use as a direct spray on birds by: (1) <b>Misting with Electric Fog Machine</b> - Mix ¾ lb. (6.4 ozs.) of this product in 1 gal. of spray. Use 1½ gals. per 1,000 Hens in cages, on litter or on slatted floors. Repeat in 4 weeks if necessary. (2) <b>Spraying with knapsack or cylinder type compressed air sprayers</b> - Mix ¼ lb. (4 ozs.) of this product in 5 gals. of spray. Use 1 gal. per 100 Hens in cage, on litter or on slatted floors. Repeat in 4 weeks if necessary. For Northern fowl mite control, ensure pressured spray penetrates feathers of vent area on birds. Direct spray misting for Chicken mites and Fleas is a supplement to spraying rooms and buildings for control of these pests.		
Poultry quarters	Bedbugs, Chicken mites, Fleas, Lice, Northern fowl mites	5 (4½ tbsps. per gal.)
	SPECIFIC DIRECTIONS: Apply 1 to 2 gals. of spray mixture with conventional power spray or knapsack equipment per 1,000 sq. ft. of wall, litter or roost surface. Force spray into cracks. Repeat as needed. Avoid spraying nests, eggs and feeding and water troughs.	
	Fowl ticks	20 (17 tbsps. per gal.)
SPECIFIC DIRECTIONS: Ventilate while spraying.		

**POULTRY INSECT CONTROL**

Darkling beetles (Lesser mealworms or Litter beetles)

Site	Pest	Lbs. of This Product
Poultry quarters	Darkling beetles (Lesser mealworms or Litter beetles)	0.625 to 1.25 lbs. per 1,000 sq. ft.
	<b>SPECIFIC DIRECTIONS:</b> Apply as a dilute spray to floor surface, walls, cracks, posts and crevices with sufficient volume of finished spray to wet all surfaces thoroughly or apply as a dry dust to floor surface, walls, cracks, posts and crevices using rotary, mechanical or electrostatic duster. Use face mask and avoid contact with dust. Optimum application timing is immediately after removing old flock from, or before placing new flock into the house when Beetles are most active. Ventilate during application. Avoid contamination of feeding and watering units.	
<b>RESTRICTIONS AND PRECAUTIONS:</b> Do not treat Birds or premises within 7 days of slaughter.		

**WARRANTY—CONDITION OF SALE**

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool, dry area away from any heat or ignition source. Do not allow storage in a damp or humid area. Do not stack over 2 pallets high. Move bags carefully so as not to tear or puncture. Do not move from one area to another unless they are securely sealed to prevent dust from escaping. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids and alkalis. Store in original containers only. If the contents are leaking or material is spilled, follow these steps while wearing protective equipment.

1. Collect and place in suitable containers for disposal.
2. Wash area with soap and water to remove remaining pesticides.
3. Follow washing with clear water rinse.
4. Do not allow runoff to enter sewer or contaminate water supplies.
5. Dispose of waste as indicated below:

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.