FIFRA Section 24(c) Special Local Need Label



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA



EPA Reg. No. 62719-282 EPA SLN OR-110007

(For Distribution and Use Only Within the State of Oregon)

Control of Spotted Wing Drosophila in Cherry

Expiration Date: This label is valid until December 31, 2021 or until otherwise amended, withdrawn, canceled, or suspended.

Group	5	INSECTICIDE
Active Ingredient: spinosad (a mixture of spinosyn A and spinosyn D)80% Other Ingredients20%		

Contains 80% active ingredient on a weight basis.

CAUTION

Environmental Hazards

This product is toxic to bees exposed to treatment for 3 hours following treatment. Do not apply this pesticide to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on the label.

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Special Local Need (SLN) and the EPA registered Entrust product label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Refer to product label for Entrust for General Use Precautions, Mixing Directions, and Application Directions. Follow all use precautions and restrictions specified on the label.

Application Timing: Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply 1.25 to 2 oz of Entrust per acre as a foliar spray to control spotted wing drosophila. Use a higher rate in the rate range for larger larvae or moderate to severe pest infestations. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: Rotate to a different class of insect control products after two applications of Entrust. Do not make more than three applications per acre per year of Group 5 insecticides (spinetoram and spinosad).

Specific Use Restrictions:

- **Pre-Harvest Interval**: Do not apply within 3 days of harvest.
- Treatment Schedule: If three applications are made, one may be made up to 30 days PHI, another up to 10 days PHI, and the final application at 3 days PHI. If two applications are made, the time interval between applications is restricted to at least 7 days.
- Do not apply more than 2 oz of Entrust (0.1 lb ai) per acre per application.
- Do not apply more than a total of 6 oz of Entrust (0.3 lb ai) per acre per year.
- Do not apply more than a total of 4 oz of Entrust per acre within 10 days of harvest.
- Do not make more than three foliar applications per acre per year of Group 5 insecticides (spinetoram and spinosad). This restriction supersedes the number of applications on the label affixed to the container of Entrust.

24(c) Registrant: Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, Indiana 46268 1-800-992-5994 (option #4)

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow R184-065 Accepted: 12/29/16 Replaces R184-049.