2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONTTM LANNATE® SP INSECTICIDE EPA Reg. No. 352-342

USE OF FOLIAR APPLICATION OF LANNATE® SP FOR CONTROL OF BAGRADA BUG IN BROCCOLI, CABBAGE, CAULIFLOWER, AND CHINESE CABBAGE IN THE STATES OF ARIZONA AND CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

RESTRICTED USE PESTICIDE DUE TO HIGH ACUTE TOXICITY TO HUMANS

For retail sale use only by Certified Applicators or persons under their direct supervision and Only for those covered by Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading or repair of application equipment.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPontTM Lannate® SP can be applied by foliar application at 0.75 to 1 pound product per acre when bagrada bug populations reach locally determined economic threshold.

Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for your area.

Crop	Rate DuPont™ Lannate® SP Lbs. Per Acre	Last Application Days to Harvest (PHI)	Pounds Lannate® SP per acre/crop	REI
Broccoli	0.75 to 1.0 lb.	3	No more than 7 Lbs.	48 hrs.
Cabbage	0.75 to 1.0 lb.	1	No more than 8 Lbs.	48 hrs.
Cauliflower	0.75 to 1.0 lb.	3	No more than 8 Lbs.	48 hrs.
Chinese				
Cabbage	0.75 to 1.0 lb.	10	No more than 8 Lbs.	48 hrs.

APPLICATION

Application can be made by ground or air application only. Application of DuPontTM Lannate® SP should be repeated as necessary to prevent reinfestation from adults migrating into the crop. For broccoli, cabbage and cauliflower the minimum application interval is 2 days and for Chinese cabbage the minimum application interval is 5 days. Use sufficient water to obtain thorough, uniform coverage. Since Lannate® is a fast acting contact insecticide, best results follow direct spraying of target insects. Bagrada bugs are very mobile pests. They may reinfect the treated area quickly.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT or at http://cropprotection.dupont.com/

© 2012 E.I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898