

**SUPPLEMENTAL LABELING**

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

**Syngenta Crop Protection, LLC**

P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1431A-S15 1211**

**Gramoxone® SL 2.0**

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide  
Vegetables (Cucumber, Melons, Pumpkin, Squash) – For Use Only in the State of Hawaii

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) ..... 30.1%

Other Ingredients: ..... 69.9%

Total: ..... 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.  
Contains alerting agent (odor), emetic, dye.

**EPA Reg. No. 100-1431**

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON  
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone SL 2.0 Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<p><b>VEGETABLES</b></p> <p>Cucumber Melons Pumpkin Squash</p> <p>For Use Only in the State of Hawaii</p>	3	Directed/ Shielded Spray	1.9 pts.	Ground: 20 gals.	8 days	<ul style="list-style-type: none"> <li>• For control or suppression of emerged weeds between rows after crop establishment.</li> <li>• Use properly configured precision spray equipment to avoid contact of spray with the crop.</li> <li>• Apply when weeds are 6" or less in height and actively growing.</li> <li>• Always use a high quality nonionic surfactant or a crop oil concentrate.</li> <li>• Use even flat fan tips at 20 to 30 psi pressure at the nozzle, or use twin even flat spray or hollow cone spray tips at 30 psi pressure at the nozzle. Do not use tips that require higher operating pressures making spray droplets susceptible to drift. Do not use tips that produce large droplets as performance will be reduced.</li> <li>• Do not allow spray to contact green stems or foliage, as injury may result.</li> <li>• Avoid excessive speed as it may cause spray to contact the crop when the boom bounces.</li> <li>• Do not spray in high wind favoring drift into the crop.</li> <li>• Do not use on soils with less than 5% clay.</li> <li>• Do not apply after first bloom and within eight (8) days of harvest.</li> <li>• Do not harvest within eight (8) days of application.</li> <li>• Do not allow animals to graze in treated areas.</li> <li>• DO NOT apply this product through any type of irrigation system.</li> </ul>

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S15 1211 – tb – 02/08/12