

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Syngenta Crop Protection, LLC

P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1431A-S1 1211

Gramoxone® SL 2.0

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide
CITRUS HYBRIDS (Tangelos, Tangors, Temple orange, Clementines)
For Use Only in the State of California

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride)..... 30.1%

Other Ingredients: 69.9%

Total: 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.

Contains alerting agent (odor), emetic, dye.

EPA Reg. No. 100-1431

KEEP OUT OF REACH OF CHILDREN.



**DANGER / POISON
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone SL 2.0 Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<p>CITRUS HYBRIDS</p> <p>(Tangelos, Tangors, Temple orange, Clementines)</p> <p>For Use Only in the State of CALIFORNIA</p>	5	Directed Spray	2.5 - 4.0 pts.	Ground: 10 gals.	-	<ul style="list-style-type: none"> • Maximum of 5 applications per year. • Pre-harvest Interval – 0 days. • DO NOT allow spray to contact green stems (except suckers), fruit or foliage. • Use a shield or wrap plant when spraying around young trees. • DO NOT graze treated areas. • DO NOT feed cover crops grown in treated areas to livestock. • DO NOT apply when weather conditions favor drift from treated areas. • DO NOT apply by ground equipment within 25 ft. or by air within 75 ft. of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries; and commercial fish ponds. • Make ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph for ground applications or 7 mph for aerial applications. • Follow recommended spray volumes listed. These are minimum volumes only, and spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage. • DO NOT apply this product through any type of irrigation system. • DO NOT enter treated field within 12 hours of application.

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S1 1211 – tb – 02/08/12