

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F

EPA Reg. No. 62719-442

EPA 24(c) Special Local Need Registration SLN CA-060026
For Distribution and Use Only in the State of California

Control of Western Yellowstriped Armyworm Infesting Sweet Potato

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid[®] 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Application Rates, Application Timing and Restrictions:

Application Rate (fl oz/acre)	Application Timing	Restrictions
6 - 10 (0.09 – 0.16 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult.	<ul style="list-style-type: none">• Do not apply more than 10 fl oz of Intrepid 2F (0.16 lb ai) per acre per season.• Do not make more than 1 application per acre per season.• Pre-harvest Interval: Do not harvest within 7 days of application.• See Rotational Crop Restrictions below.

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment. Higher spray volumes may be required to provide thorough coverage to a larger, more mature crop. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-043

Accepted: 10/23/06

Initial printing.