

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F

EPA Reg. No. 62719-442

EPA 24(c) Special Local Need Registration SLN NC-060003
For Distribution and Use Only in the State of North Carolina

Control of Lepidoptera Larvae Infesting Sweet Potato

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- All applicable directions, restrictions, and precautions on the registered product label for Intrepid[®] 2F insecticide (Dow AgroSciences EPA Reg. No. 62719-442, Rohm and Haas EPA Reg. No. 707-277) are to be followed. Read the label affixed to the container for before applying.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworms cabbageworms loopers saltmarsh caterpillar webworms	6 - 10 (0.09 – 0.16 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> • Do not apply more than 30 fl oz of Intrepid 2F (0.48 lb ai) per acre per season. • Do not make more than 3 applications per acre per season. • Pre-harvest Interval: Do not harvest within 7 days of application. • See Rotational Crop Restrictions below.

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment. Higher spray volumes may be required to provide thorough coverage to a larger, more mature crop. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-049

Accepted: 07/18/06

R846-046