

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX® HERBICIDE
BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN
THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



EPA Reg. No. 524-537

Roundup WeatherMAX and TranSorb are
trademarks of Monsanto Technology LLC

FOR CONTROL OF ANNUAL AND PERENNIAL WEEDS IN SUNFLOWER AND SAFFLOWER

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

DIRECTIONS FOR USE

LABELED CROPS: Safflower, Sunflower.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, At-Planting, Hooded Sprayers in Row Middles, Shielded Sprayers in Row Middles, Wiper Applications in Row Middles, Preharvest, and Post-Harvest treatments.

NOTE: Unless otherwise specified, weed control applications may be made according to the rates listed in the "ANNUAL WEEDS", and "PERENNIAL WEEDS RATE SECTION" in the Roundup WeatherMAX herbicide booklet label.

Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting sunflower or safflower crops. Broadcast applications must be made prior to emergence.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl® may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue.

PRECAUTIONS, RESTRICTIONS: In safflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 2 quarts of this product per acre per year. In sunflower, the total combined application rate of all preemergence and hooded/shielded

sprayer treatments must not exceed 22 fluid ounces of this product per acre per year.

Hooded or Shielded Sprayers in Row Middles, Wiper Applications in Row Middles

USE INSTRUCTIONS: Wiper applicators or hooded/shielded sprayers may be used between the rows once the crop is established. See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the Roundup WeatherMAX herbicide label for information on proper use and calibration of this equipment.

PRECAUTIONS, RESTRICTIONS: In safflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 2 quarts of this product per acre per year. In sunflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 22 fluid ounces of this product per acre per year.

Preharvest

USE INSTRUCTIONS: This product provides weed control when applied as a harvest aid to a physiologically mature crop prior to harvest of sunflower or safflower. For safflower, apply when seed has lost its opaque character, approximately 20 to 30 days after the end of flowering of the secondary branches. For sunflower, apply when the backsides of sunflower heads are yellow and bracts are turning brown and seed moisture content is less than 35%.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or livestock feeding. Apply no more than 2 quarts of this product at a preharvest timing to safflower. Apply no more than 22 fluid ounces of this product at a preharvest timing to sunflower. Applications must be made at least 30 days prior to planting any crop not listed on the Roundup WeatherMAX herbicide label booklet.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of safflower or sunflower. Higher rates may be required for control of large weeds, which were growing in the crops at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. Applications must be made at least 30 days prior to planting any crop not listed on the Roundup WeatherMAX herbicide label booklet.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

All other trademarks are the property of their respective owners.

©2007 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

1/9/2006

63003F1-61