



**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA**

**Caparol® 4L**

**EPA Reg. No. 100-620  
EPA SLN No. CA-980017**

**For Control Of Certain Weeds In Transplanted Celery**

Active Ingredient:

Prometryn: 2,4-bis(isopropylamino)-6-(methylthio)-s-triazine .....	44.4%
Other Ingredients .....	55.6%
Total: .....	100.0%

Caparol 4L contains 4 lbs. active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

---

**DIRECTIONS FOR USE**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Caparol 4L label.

This label must be in the possession of the user at the time of pesticide application.

To amend Caparol 4L label to include pre-transplant applications:

**Single Pre-Transplant Application**

On coarse-textured soils and soils low in organic matter, make one application of 2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water. On fine-textured soils and soils high in organic matter, make one application of 3.2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water. Make the application up to 21 days prior to transplanting. Make the application to a weed-free bed and do not disturb the treated area between application and

transplanting. Note: A post-transplant application will not be allowed if the pre-transplant application was made at the maximum allowable use rate (4 pts. per acre).

### **Split Pre-Transplant and Post-Transplant Application:**

On coarse-textured soils and soils low in organic matter, the lower rate range of 2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water is recommended. The maximum post-transplant application allowable on coarse-textured soils and soils low in organic matter is 3.2 pts. per acre.

On fine-textured soils and soils high in organic matter, the higher rate range of 3.2 to 4 pts. per acre of Caparol 4L in a minimum of 20 gallons of water is recommended.

Make the pre-transplant application up to 21 days prior to transplanting. Make the application to a weed-free bed and do not disturb the treated area between application and transplanting. A second application of Caparol 4L may be made during the 2 to 6 week period after transplanting and before weeds are two inches tall. The total amount of Caparol 4L applied to the crop (pre-transplant plus post-transplant applications) cannot exceed the maximum allowable use rate (4 pts. per acre).

### **Post Transplant Application**

As indicated on the EPA-registered Caparol 4L label make one application of 2 to 3.2 pts. per acre of Caparol 4L on coarse-textured soils and soils low in organic matter in a minimum of 20 gallons of water. Make one application of 3.2 to 4 pts. per acre of Caparol 4L on fine-textured soils and soils high in organic matter in a minimum of 20 gallons of water. Application may be made over-the-top to celery during the 2 to 6 week period after transplanting and before weeds are two inches tall.

**NOTE:** To avoid possible illegal residues, do not make a Caparol 4L application within 45 days of harvest.

**Rotational Crops:** The following crops may be seeded five months after applying no more than 4 pts. of Caparol 4L per acre on celery: cabbage, celery, corn, onions, peas, and red beets.

Caparol® 4L is a trademark of Syngenta

24(c) registrant:  
Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, NC 27419-8300

EPA SLN No: CA-980017

Label Code: CA0620052AA0204