Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Starane[®] NXT

EPA Reg. No. 62719-557

Special 2(ee) Recommendation[†]
(For Distribution and Use in the States of Colorado, Kansas, and Nebraska)

For Postemergence Control of Annual Broadleaf Weeds In Sorghum (Grain and Forage)

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Starane® NXT insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Starane NXT according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Starane NXT.

Directions for Use

Refer to product label for Starane® NXT for General Use Precautions, Mixing Directions and Application Directions.

Apply 21 fl oz per acre of Starane NXT at the following crop stages for selective control of annual broadleaf weeds in grain and forage sorghum:

- Prior to planting to just prior to crop emergence
- 4-leaf stage through the 7-leaf stage
- 8-leaf stage to prior to the boot stage using drop nozzles

Refer to the label affixed to the container for Starane NXT for resistance management recommendations, all use directions, precautions, and restrictions.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R052-013 Issued: 07/29/13 Initial printing.