



ATRA

Section 24(c) Special Local Need Label

**RESTRICTED USE PESTICIDE
(GROUND AND SURFACE WATER CONCERNS)**

**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED
APPLICATOR'S OR PERSONS UNDER THEIR DIRECT
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE
CERTIFIED APPLICATOR'S CERTIFICATION.**

**THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO
GROUND AND SURFACE WATER CONCERNS. USERS MUST READ
AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS
FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE
TO REACH GROUND AND SURFACE WATER.**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
OKLAHOMA**

AATREX[®] NINE-O[®]

**EPA Reg. No. 100-585
EPA SLN No. OK-830030**

**For Control Of Broadleaf And Grass Weeds Following Wheat Harvest
In A Wheat-Fallow-Wheat Rotation Using AAtrex Nine-O Alone Or In
Tank Mixtures.**

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA registered AAtrex Nine-O label and on the labels of herbicides tank mixed with AAtrex.

This labeling must be in the possession of the user at the time of pesticide application.

DO NOT contaminate or apply to ponds, lakes, or streams.

AAtrex is a broad spectrum herbicide for preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat, during the period following wheat harvest in a winter wheat-fallow-winter wheat rotation.

For best results, apply herbicide after straw has been uniformly distributed over the wheat stubble or soil surface.

Winter Wheat - For Weed Control After Harvest in a Winter Wheat – Fallow - Winter Wheat Rotation.

Precautions: **DO NOT** apply on soils having a pH greater than 7. **DO NOT** apply in the Oklahoma Panhandle. **DO NOT** treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil. **DO NOT** treat soils with calcareous surface layers. **DO NOT** use on sands. Avoid spray overlap.

Apply AAtrex Nine-O at 1.1 lb./A as soon as possible following wheat harvest.

For control of established weeds, tank mix with a contact herbicide registered for this use, such as Gramoxone Inteon™ at 1.5-3.0 pts./A + X-77® at 1-2 pts. per 100 gallons of spray solution, or stubble plow to control large established weeds. Refer to the herbicide labels for additional directions, weeds controlled, precautions, and limitations.

Apply in a minimum of 20 gals. of water per acre with ground equipment.

If weeds or volunteer wheat become a problem, a shallow cultivation or a postemergence application of a registered herbicide such as Gramoxone Inteon + X-77 should be employed.

Tillage and/or application of registered herbicides will be necessary for weed control the spring following AAtrex application.

Note: Treat only once during the same fallow period. **DO NOT** plant wheat until 15 months after AAtrex application.

Chemigation: **DO NOT** apply these products through any type of irrigation system.

AAtrex®, Nine-O®, and Gramoxone Inteon™ trademarks of a Syngenta Group Company
X-77® trademark of Loveland Industries, Inc.

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: OK-830030

OK0585030CA1007