
**RESTRICTED USE PESTICIDE
Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision,
and only for those uses covered by the Certified Applicator's certification.

INTRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA

GRAMOXONE® SL 2.0

**EPA REG. NO. 100-1431
EPA SLN NO. GA-120004**

**For Suppression and/or Control of Palmer Amaranth and Florida Beggarweed
in Peanut**

KEEP OUT OF REACH OF CHILDREN

Danger/Peligro
POISON



This label expires and must not be distributed or used in accordance with this SLN registration
after December 31, 2018

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with all applicable
directions, restrictions, and precautions found in the labeling of the federally registered product
and accompanying supplemental labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to
the Worker Protection Standards on the EPA-registered Gramoxone SL 2.0 label.

This label must be in the possession of the user at the time of application.

Application Method

Gramoxone SL 2.0 can be applied through recirculating rope wick or carpet roller equipment for suppression or control of glyphosate-resistant Palmer amaranth and Florida beggarweed or to prevent seed production.

Make application at least 6 inches above the peanut canopy, and set application equipment to avoid dripping. Low ground speed enhances coverage and hence weed control. Speeds less than 5 mph are recommended.

Control of large weeds is enhanced if application is made in late afternoon or early evening.

Follow application equipment manufacturer's instructions.

Mixing Instructions and Application Rate

Mix 1 part of Gramoxone SL 2.0 with 1-1.5 parts of water to prepare a 40-50% solution. Add nonionic surfactant containing 75% or more surface-active agent at a minimum of 0.25% v/v (2 pt/100 gal or 0.32 fl oz/gal) of the finished volume.

Adjust equipment to apply up to 2 pt/A of the herbicide-water mixture .

Restrictions

1. Do not apply more than 1 pt/A of Gramoxone SL 2.0 using the recirculating rope wick or carpet roller application method.
2. Do not exceed 0.5 lb ai/A (1 qt/A) total for the entire growing season from all applications on this or any other supplemental or Section 3 Federal label for peanut
3. Make applications of Gramoxone SL 2.0 as early as possible, and not less than 30 days before harvest to control weeds that may interfere with harvesting operations.
4. Do not allow livestock to graze in treated areas.
5. Do not feed hay or threshings from treated fields to livestock.

Section 24(c) registrant:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

Gramoxone® trademark of a Syngenta Group Company

GA1431045CA1013