

## MORRA

## Section 2 (ee) Recommendation

Date: February 19, 2014

Product: Omega® 500F

EPA Reg. No. 71512-1-100

Use: Addition of Downy Mildew (*Phytophthora phaseoli*) as a Pest on Lima Bean

State: DE, MD, NJ, PA, TN, VA

## **Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Pest Controlled	Rate	Application Interval/ Timing	Instructions
Downy Mildew (Phytophthora phaseoli)	0.5 to 0.85 pints per acre	7-10 days	Application Directions: For control of downy mildew, make the first application at 10-30% bloom (i.e., when 10-30% of the plants have at least one (1) open bloom). If needed, a second application may be applied 7 to 10 days later. Use adequate water to provide coverage of foliage and flowers. Under conditions favorable for severe disease development, use the 0.85 pint rate.  Restrictions: DO NOT use more than 1.75 pints per acre per growing season. DO NOT apply within 14 days of harvest for edible-podded and succulent beans (14-day PHI). DO NOT apply within 30 days of harvest for dry and lima beans (30-day PHI). Restricted Entry Interval, REI = 12 hours.  Omega 500F may be applied through sprinkler system irrigation equipment on beans. See calibration directions on the Omega 500F Section 3 label.

The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.

The user should have this recommendation in its possession at the time of use.

This recommendation for use of Omego 500F is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

2(ee) Registrant: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419-8300

Omega® is a registered trademark of Ishihara Kaisha, LTD; Syngenta logo is a trademark of a Syngenta Group Company

©2014 Syngenta

Expiration Date: February 19, 2018

Label Code: DE0001029AC0214