

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F

EPA Reg. No. 62719-442

Control of Carob Moths Infesting Dates (Not for Use and Distribution in New York State)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid[®] 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Target Pest	Intrepid 2F fl oz/acre	Application Timing	Restrictions
carob moth	10 - 20 (0.16 – 0.31 lb ai/acre)	<p>For control of light to moderate infestations, begin applications before egg hatch of each generation and before the larvae penetrate the fruit. Once applied, the product provides 10 to 18 days of protection depending upon application rate and how rapidly fruit is expanding. Consult local spray timing advisories or follow biofix dates based on pheromone trap catches to time sprays appropriately.</p> <p>For continuous moth flight and egg laying, use the highest labeled rate. Maintain coverage on the fruit surface with 10- to 18-day re-treatment intervals.</p> <p>Alternate or intersperse with other insecticides with different modes of action targeted for the same pest so long as the re-treatment interval does not exceed the period of effectiveness of the products being alternated and Intrepid 2F is applied before larvae penetrate the fruit.</p>	<ul style="list-style-type: none">• Do not apply more than 20 fl oz per acre per application or a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per season.• Preharvest Interval: Do not apply within 7 days of harvest.• Minimum Re-treatment Interval: 10 days

Ground Application: Apply a minimum of 100 gallons per acre. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Spray Adjuvants: The addition of agricultural adjuvants to sprays of Intrepid 2F can improve initial spray deposits, redistribution and weatherability, and is highly recommended for improved efficacy. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-066

EPA accepted 10/29/13

Initial printing.