

Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Dithane[®] DF Rainshield[™]

EPA Reg. No. 62719-402

EPA 24(c) SLN No.: WA-020028

(For Distribution and Use Only Within Washington State)

For Disease Control in Small Seeded Vegetable Seed Crops

Expiration date: This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2018.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the product container before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label attached to the product container.

Directions for Use

Begin applications of Dithane[®] DF Rainshield[™] fungicide when disease first appears or is reported in the area. Apply Dithane DF Rainshield fungicide at the rate of 2 pounds per acre in thorough coverage sprays. Repeat applications on a 7 to 10 day application schedule until the threat of disease is over.

Use sufficient spray volume to achieve thorough coverage. See package label for use directions and restrictions for ground, aerial, and/or chemigation applications.

Dithane DF Rainshield fungicide may be used on the following seed crops for control of listed diseases:

Seed Crop	Diseases Controlled
Allium spp. (Leek, bunching and dry bulb onions)	Botrytis leaf blight (<i>Botrytis</i> spp.) Downy mildew (<i>Peronospora destructor</i>) Neck rot (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>)
Arugula	Alternaria leaf spot (<i>Alternaria</i> sp.)
Beet	Alternaria leaf spot (<i>Alternaria</i> sp.) Cercospora leaf spot (<i>Cercospora beticola</i>) Downy mildew (<i>Peronospora farinosa</i>) Rust (<i>Uromyces betae</i>)
Carrot	Leaf blight (<i>Alternaria dauci</i>) Leaf spot (<i>Cercospora carotae</i>) Rust (<i>Uromyces betae</i>)
Coriander	Leaf spot (<i>Alternaria</i> spp.)
Brassica and Raphanus spp.: Broccoli raab, Brussels sprout, cabbage, cauliflower, Chinese cabbage, Chinese mustard, collard, kale, kohlrabi, rutabaga, turnip, mustard,	Black spot (<i>Alternaria alternata</i> and <i>A. brassicola</i>) Cercospora leaf spot (<i>Cercospora brassicicola</i>) Downy mildew (<i>Peronospora parasitica</i>)

radish, rape, spinach mustard	
Dill	Alternaria leaf spot (<i>Alternaria</i> sp.)
Endive	Downy mildew (<i>Bremia lactucae</i>) Leaf spot (<i>Cercospora cichoria</i>)
Lettuce	Anthrachnose (<i>Microdochium panattoniamnum</i>) Cercospora leaf spot (<i>Cercospora longissima</i>) Downy mildew (<i>Bremia lactucae</i>)
Parsley	Alternaria leaf spot (<i>Alternaria</i> sp.)
Parsnip	Alternaria leaf spot (<i>Alternaria</i> sp.)
Spinach	Downy mildew (<i>Peronospora effusa</i> and <i>P. farinosa</i>) Leaf spot (<i>Alternaria</i> spp.) Leaf spot (<i>Cercospora beticola</i>) Leaf spot (<i>Cladosporium variable</i>)
Swiss Chard	Cercospora leaf spot (<i>Cercospora beticola</i>) Rust (<i>Uromyces batae</i>)

Restrictions/Precautions

- Do not apply more than 25.6 lb (19.2 lb active ingredient) per acre per crop.
- All leek, bunching and dry bulb onions, arugula, beet, carrot, coriander, broccoli raab, brussels sprout, cabbage, cauliflower, Chinese cabbage, Chinese mustard, collard, kale, kohlrabi, rutabaga, turnip, mustard, radish, rape, spinach mustard, dill, endive, lettuce, parsley, parsnip, spinach, and Swiss chard seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
- No portion of the leek, bunching and dry bulb onions, arugula, beet, carrot, coriander, broccoli raab, brussels sprout, cabbage, cauliflower, Chinese cabbage, Chinese mustard, collard, kale, kohlrabi, rutabaga, turnip, mustard, radish, rape, spinach mustard, dill, endive, lettuce, parsley, parsnip, spinach, and Swiss chard seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Leek, bunching and dry bulb onions, arugula, beet, carrot, coriander, broccoli raab, brussels sprout, cabbage, cauliflower, Chinese cabbage, Chinese mustard, collard, kale, kohlrabi, rutabaga, turnip, mustard, radish, rape, spinach mustard, dill, endive, lettuce, parsley, parsnip, spinach, and Swiss chard seed shall bear a tag or container label which forbids use of the seed for human consumption or animal
- Leek, bunching and dry bulb onions, arugula, beet, carrot, coriander, broccoli raab, brussels sprout, cabbage, cauliflower, Chinese cabbage, Chinese mustard, collard, kale, kohlrabi, rutabaga, turnip, mustard, radish, rape, spinach mustard, dill, endive, lettuce, parsley, parsnip, spinach, and Swiss chard seed may not be distributed for human consumption or animal feed.
- Applicants that wish to apply for an SLN registration under the nonfood/nonfeed designation must obtain a copy of WSDA seed crop rules and consult with WSDA.
- **Chemigation:** For use under this SLN label do not apply this product through any type of irrigation system.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

WSDA Aquatic Advisory

This product is toxic to fish. Dithane DF Rainshield should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the Washington State Department of Agriculture Endangered Species web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx>, or contact the Washington Department of

Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Washington State Department of Agriculture Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the Washington State Department of Agriculture Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in the state of Washington.

24c Registrant:

**Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054 USA**

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R818-066

Accepted: 04/09/14

Replaces R818-038