



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Leverage® 360 Insecticide

For Use on Grape

EPA Reg. No. 264-1104

Product Bulletin

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

GRAPE

Includes: Table grape, Raisin, Wine and Muscadine grape

Pest Controlled	Rate fluid ounces/Acre
Japanese beetle (adult)	3.2 – 6.4

RESTRICTIONS

Pre-Harvest Interval (PHI): **3 day**

Minimum interval between applications: **14 days**

Maximum LEVERAGE 360 Insecticide allowed per crop season: **6.4 fluid ounces/A** (0.05 lb AI/Acre β -cyfluthrin and 0.10 lb AI/Acre imidacloprid)

NOTES

LEVERAGE 360 Insecticide is to be applied through properly calibrated ground application equipment, only.

Minimum application volume (water): 25 GPA – ground, 25 GPA – aerial application.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)
or

Visit our worldwide web site at www.bayercropscienceus.com

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

Leverage is a registered trademark of Bayer.