RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) Leverage[®] 360 Insecticide EPA Reg. No. 264-1104

For control of Cabbage Maggot, Garden Symphylan and Seed Corn Maggot in Brassica Leafy Vegetables Crop Group 5 and Leafy Greens Crop Group 4A

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

BRASSICA LEAFY VEGETABLES

Crops of Crop Group 5, plus Turnip greens, including: Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccolo, Chinese broccoli (*gai lon*), Chinese cabbage (*bok choy*), Chinese cabbage (*napa*), Chinese mustard cabbage (*gai choy*), Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, Turnip greens.

LEAFY GREENS

Crops of Crop Subgroup 4A Including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach [including New Zealand and vine (Malabar spinach, Indian spinach)]

PESTS CONTROLLED	Rate Leverage 360 Insecticide fluid ounces/Acre	Method
Cabbage Maggot Garden Symphylan Seed Corn Maggot	3.0 fl.oz. per acre	For reduction of damage caused by the listed pests, apply specified dosage per acre to the soil as a soil-incorporated treatment before or at planting or transplanting.

Notes and Restrictions:

Maximum LEVERAGE 360 Insecticide allowed per crop season: 12.8 fluid ounces/A (0.1 lb Al/A β-cyfluthrin, 0.2 lb Al/A imidacloprid)

Maximum β -cyfluthrin allowed in all forms per crop season: 0.1 lb Al/A Maximum cyfluthrin allowed in all forms per crop season: 0.2 lb Al/A

Maximum β -cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.2 lb Al/A Maximum imidacloprid allowed in all forms as a soil application per crop season: 0.38 lb Al/A

Not for use on crops grown for seed.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937)