



Bumper ES
EPA REG. NO. 66222-241

FIFRA §2(ee) RECOMMENDATION
For control of Rust and Scab on Almonds in California Only.

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by U.S. EPA.

APPLICATION INFORMATION

CROP	PESTS CONTROLLED	RATE OF APPLICATION PER ACRE	APPLICATION INSTRUCTIONS
Almonds	Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>)	6.4 to 8 fl oz	Apply 6.4 to 8 fl oz (0.18 – 0.225 lbs. active) per acre. Begin application approximately 2 – 5 weeks after petal fall when environmental conditions are favorable for disease development. For optimum disease control, apply on a 7- to 14- day application schedule. Make no more than two consecutive applications before rotating to a fungicide with a different mode of action (not a Group 3).
	<ul style="list-style-type: none"> • Do not apply more than 32 fl oz per acre per season of Bumper ES. • Do not apply more than 0.90 lb a.i. Propiconazole per acre per season • Do not apply within 60 days of harvest. • Do not graze livestock in treated areas or cut treated cover crop for feed. 		

Read the “**LIMITATION OF WARRANTY AND LIABILITY**” in the label booklet for Bumper ES before using.

060512
EPA SAL DATE 043012



3120 Highwoods Blvd, Suite 100, Raleigh NC 27604
T: 919.256.9305 ♦ F: 919.256.9308