CORE 6% MANGANESE EDTA

For correction or prevention of manganese deficiency in agricultural crops, and vegetable crops

GENERAL INFORMATION

Manganese EDTA is a fully chelated, plant available, liquid micronutrient and ethylenediaminetetraacetic acid chelating agent for use in vegetable crops, field crops, trees, and vines. Manganese EDTA when used along with a sound fertility program assures the nutritional requirements of the plant.

Core 6% Manganese EDTA is a liquid micronutrient compatible with liquid fertilizers containing polyphosates and othophosphates. This product is a stable manganese chelate and is both ammoniacal nitrogen and sodium free which provides a low odor formulation.



GUARANTEED ANALYSIS:

Soluble Potash (K,0)	10.0%
Manganese (Mn)	6.0%
6.0% Chelated Manganese	

Derived from: Potassium Hydroxide and Manganese EDTA

Function of Manganese in Plants

Manganese serves as a activator for enzymes in the plant growth processes. It assists iron in chlorophyll formation. Manganese is necessary in photosynthesis, N metabolism and to form other components required for plant metabolism. It is an essential component of some enzyme systems.

There are a number of functions or processes that manganese is part of in plants including:

- Functions in the formation of riboflavin, ascorbic acid, and carotine.
- Essential for assimilation of carbon dioxide in photosynthesis.
- Serves as a catalyst for enzyme reaction.
- Aids in synthesis of chlorophyll and functions in photosynthesis and nitrate assimilation
- Serves in oxidation-reduction reactions such as to change Fe +++ to Fe++ in the plant

Manganese Deficiency in Plants:

- Manganese deficiencies mainly occur on organic soils, high-pH soils, sandy soils low in organic matter and over limed soils.
- Interveinal chlorosis of young leaves
- Gradation of pale green leaf coloration, with darker color next to veins.
 There is no sharp color distinction between veins and interveinal areas as there is with iron deficiency.
- Development of grey speks (oats), interveinal white streaks (wheat), interveinal brown spots or streaks (barley), or necrotic spots (potatoes).
- Some crops which are sensitive to manganese deficiency are: corn, cotton, oats, potatoes, soybean, sugar beets, wheat, dry bean, cauliflower, celery, lettuce, onion, spinach, citrus, and peach.

0-0-10



Product Specs

Product class Fertilizer
Chemical Formula K₂Mn EDTA
CAS: 68013-77-0
Appearance Clear to slight pink
Specific gravity 1.32

Solubility in water miscible
Odor no odor
Formulation liquid
Molecular Weight 421.4
Weight/gallon 11 lbs at 68°

pH (5% solution) 7.0 Appearance clear/pink Min. storage temp. 32°F

Package Sizes: Bulk, 250 gallon mini-bulks, 30 gal drums, 5 gal pails, 2 x 2.5 gallon package

Product is available from Hanford, CA and Humboldt. TN

This bulletin provides some technical information and is not intended to give complete information for all applications.

Always read and follow label directions.



FOR MORE INFORMATION, LABELS, MSDS and a COMPLETE PRODUCT LIST:

log onto www.CoreAgri.com
contact your sales representative
or call: 1-800-385-4715

COREAGRI, LLC

P.O. Box 1027 • Arroyo Grande, CA 93420 • USA customerservice@coreagri.com | 800-385-4715