



CORE Zinc Nitrate

APPLICATION INFORMATION



CORE Zinc Nitrate can be used on any agricultural crop where a zinc deficiency exists or where added zinc would be beneficial. Compatibility: **CORE Zinc Nitrate** is formulated to mix with most commonly used insecticides and fungicides. This product can also be applied via irrigation. However, a jar test for compatibility should be performed prior to application. This product is not recommended for use with copper or tin fungicides.

GENERAL RECOMMENDATIONS – FOLIAR APPLICATIONS

Apples: Applications of 1/2 to 1 pint/acre at 1/2 inch green to pink stage. In season application can be made to the foliage 3 to 4 weeks after petal fall of 1/4 to 1/2 pint/acre. Applications 1/2 to 1 pint/acre after harvest and within two weeks of leaf fall. Water Rate: 50 to 100 gallons/acre.

Blueberries: Applications of 1/4 to 1/2 pints/acre starting from leaf expansion to sign of small fruit. Water Rate: 50 gallons per acre.

Canola: 1/2 to 1 pint/acre prior to pod formation. If appropriate, consider a second application 10 to 14 days later. Water Rate: 20 gallons/acre.

Carrots (Field Grown): 2 to 3 applications of 1/4 to 1.5 pints/acre when crop is 4 - 6 inches tall. Allow 10 to 14 day intervals between applications. Water Rate: 20 gallons/acre.

Cereals: 1/2 to 1 pint/acre from top dress to heading. Water Rate: 20 gallons/acre.

Citrus: Applications of 1/2 to 1 pint/acre, spring and fall flush is best, repeat as required.

Water Rate: 50 to 100 gallons/acre.

Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Calabrese, Cauliflower): Applications of 1/4 to 1/2 pint/acre from 10 to 14 days post thinning; repeat as required using 1 pint/acre. Water Rate: 20 gallons/acre.

Corn: Application of 1/2 to 1 pints/acre from 5th to 6th leaf up to just before tasseling.

Water Rate: 20 gallons/acre.

Cotton: 1/2 to 1 pint/acre at 6 to 8 leaf stage, repeat as required. If used with pesticides use 1 pint/acre.

Water Rate: 20 gallons/acre.

Cucurbits (Field Grown): Applications of 1/4 to 1 pint/acre commencing at first bloom, may repeat after each picking with 1 pint/acre. Water Rate: 20 gallons/acre.

English Walnuts: Applications of 1/4 to 1 pint/acre tender pink to early feather, repeat 1/3 pint/acre with first insect spray. Water Rate: 100 gallons/acre.

Lettuce (Field Grown): Applications of 1/4 to 1/2 pint/acre from 10 to 14 days post thinning. Repeat as required using 1 pint/acre. Water Rate: 50 gallons/acre.

Milo: Application of 1/2 to 1 pint/acre from 5th to 6th leaf up to boot. Water Rate: 20 gallons/acre.



www.CoreAgri.com