



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALABAMA, DELAWARE, FLORIDA, GEORGIA, LOUISIANA, MARYLAND, MISSISSIPPI, NEW JERSEY, NORTH CAROLINA, PENNSYLVANIA, SOUTH CAROLINA, TENNESSEE AND VIRGINIA

Acephate 97UP® Insecticide EPA Reg. No. 70506-8

For control of Kudzu bug

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	AMOUNT OF ACEPHATE 97UP PER ACRE	APPLICATION INSTRUCTIONS
Cotton (Foliar)	1 lb. (16 oz.)	Ground Application: Apply in 10 to 25 gallons of water per acre by conventional ground equipment to ensure thorough coverage. Aerial Application: Make applications in 3 to 10 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Do not apply more than 4.125 lbs. product (4 lbs. ai/A) per season. This includes all methods of application. Allow at least 21 days to elapse between final application and harvest. Minimum spray interval is 7 days.



Recommendation

CROP	AMOUNT OF ACEPHATE 97UP PER ACRE	APPLICATION INSTRUCTIONS
Soybeans	0.77 - 1 lb. (12.3 - 16 oz.)	Ground Application: Apply in 10 to 50 gallons per acre by conventional ground equipment to ensure thorough coverage. Aerial Application: Make applications in 5 to 10 gallons per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Do not apply more than 1.55 lbs product/A (1.5 lbs ai/A) per crop cycle. Allow at least 14 days to elapse between final application and harvest. Minimum spray interval is 7 days.
Beans and Lima Beans, Dry and Succulent forms (except succulent green beans, unless grown only for seed)	1 lb. (16 oz.)	Ground Application: Apply in 20 to 100 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop. Aerial Application: Make applications in a minimum of 2 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Minimum spray interval is 7 days. For dry beans, allow at least 14 days between application and harvest. For succulent lima beans, allow at least 1 day between application and harvest. Do not apply more than 2.1 lbs product/A (2 lbs. ai/A) per season. Use on succulent green beans as a food crop is prohibited (includes string beans, snap beans, and wax beans). Succulent green bean use is permitted ONLY if grown for seed.

Rev. 07/16/2013