

Supplemental Labeling

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF IDAHO.

WILBUR-ELLIS

SUPERIOR SPRAY OIL N.W.

EPA REG. NO. 2935-405 EPA SLN NO. ID-980010

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the label for EPA Reg. No. 2935-405.

CROPS: Hops

PEST: For the control of hop powdery mildew, (sphaerotheca humuli).

SITE: State of Idaho

RATE: Use 1-2 gallons per 100 gallons of water.

APPLICATION VOLUME:

Apply by ground equipment only. Thorough coverage is essential. Apply 50-150 gallons of dilute spray emulsion per acre depending on plant size. Use spray equipment that ensures producing small uniform spray droplets; best achieved with high spray pressure (minimum of 175 psi) and ceramic hollow cone nozzles.

FREQUENCY TIMING OF APPLICATIONS: Make first application at first sign of infection. Repeat every 7-14 days depending on level of new growth. Use shorter spray interval and/or higher rate when conditions for disease developing are more severe. Post-harvest applications can be made to reduce overwintering inoculum.

PRE-HARVEST INTERVAL: No interval required.

RESTRICTED-ENTRY INTERVAL: Do not enter or allow entry into treated areas during the restricted-entry interval (REI) of 12 hours.

COMPATIBILITY INFORMATION:

Do NOT Use propargite (Omite) during, with or following an oil spray.

MAXIMUM APPLICATION TEMPERATURE: Do not apply oil when the temperature at the time of application is above 90 degrees F or to heat/water stressed plants.

Do not apply this product through any type of irrigation system.

WILBUR-ELLIS Logo® and IDEAS TO GROW WITH® are registered trademarks of WILBUR-ELLIS Company.

This Special Local Need label for Superior Spray Oil N.W. expires on December 31, 2019.

Rev.02/24/14