



Recommendation

BIFENTURE™ 10DF

Insecticide/Miticide

EPA Reg. No. 70506-227

FIFRA SECTION 2(ee) RECOMMENDATION FOR CALIFORNIA

Additional Use Directions for Strawberries

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEST	RATE OUNCES BIFENTURE 10DF/A	SPECIFIC DIRECTIONS
Thrips Vinegar Flies (adults) <i>Drosophila spp.</i> (Including Spotted Winged Drosophila)	8.0 – 32.0	Apply when pest populations reach damaging thresholds. Repeat as necessary at 7-14 day intervals. Ground Application: Apply in a minimum of 50 gallons per acre. Air Application: Apply in a minimum of 5 gallons per acre. Do not apply more than 80 oz product (0.5 lb ai) per acre per season. Preharvest interval: 0 days. Thorough spray coverage is important to obtain optimum and extended control.

CALIFORNIA SPECIFIC REQUIREMENTS FOR STRAWBERRY HARVESTERS: Harvesters and other personnel performing tasks with all-day foliage contact in treated fields within five (5) days of application must wear a long-sleeved shirt, long pants, and shoes plus socks. Following treatment of strawberry fields at rates of Bifenture 10DF greater than 0.10 lb ai/acre, harvesters must wear gloves for five (5) days following application.

Observe all applicable precautions, restrictions and limitations on the EPA-approved label for Bifenture 10DF. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Rev. 7/20/10

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
www.upi-usa.com
1-800-438-6071