



Recommendation

BIFENTURE™ 10DF

Insecticide/Miticide

EPA Reg. No. 70506-227

FIFRA SECTION 2(ee) RECOMMENDATION FOR CALIFORNIA

Additional Use Directions for Caneberries

Including Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Raspberries

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEST	RATE OUNCES BIFENTURE 10DF/A	SPECIFIC DIRECTIONS
Thrips Vinegar Flies (adults) <i>Drosophila</i> spp. (Including Spotted Winged Drosophila) <i>Lygus</i> spp.	8.0 – 16.0	Apply by air or ground equipment using sufficient water to obtain full coverage of the canopy. Use a minimum of 10 gallons per acre by air and 50 gallons per acre by ground. One application may be made pre-bloom and a second application may be made post bloom. Do not apply more than 32 oz product (0.2 lb ai) per acre per season. Preharvest interval: 3 days.

Observe all applicable precautions, restrictions and limitations on the EPA-approved label for Bifenture 10DF. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Rev. 7/27/10