



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN MICHIGAN, NEW JERSEY, INDIANA, VIRGINIA, MAINE AND FLORIDA.

Bifenture 10DF
EPA Reg. No. 70506-227

Additional Use Directions for Blueberries (highbush and lowbush)

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

Pest	Rate Ounces Bifenture 10DF/A	Specific Directions
Thrips <i>Drosophila</i> spp. (adults & larvae) (Including Spotted Winged <i>Drosophila</i>) <i>Lygus</i> spp.	5.3-16.0	Apply in at least 2 gallons per acre by air or 10 gallons with ground equipment. Thorough coverage is essential to achieve adequate control. Do not apply more than 80 oz. product (0.5 lb ai) per acre per season. Use higher rates under heavy insect pest pressure. Pre harvest interval: 1 day.

Rev. 4/11/2012