



Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF  
NEW YORK**

**Caparol® 4L**

**EPA Reg. No. 100-620  
EPA SLN No. NY-140007**

**For Control of Certain Weeds in Carrot and Cilantro**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019

Active Ingredient:

Prometryn: 2,4-bis(isopropylamino)-6-(methylthio)-s-triazine .....	44.4%
Other Ingredients .....	55.6%
Total:.....	100.0%

Caparol 4L contains 4 lbs. active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

---

**DIRECTIONS FOR USE**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Caparol 4L label.

This label must be in the possession of the user at the time of pesticide application.

---

**CARROT**

---

Caparol 4L can be applied preemergence and or postemergence over the top to carrot. Read and follow all directions for use in carrot.

**Use Precautions for Pre and Postemergence Applications in Carrot**

Apply up to three applications of Caparol 4L at the rate of 2 - 4 pt/A per application.

Do not exceed one preemergence at up to 4 pt/A and two postemergence applications each at up to 2 pt/A or one postemergence application at up to 4 pt/A per crop cycle.

Do not exceed 8 pt/A of Caparol 4L per crop cycle

Make postemergence applications through the 6 leaf stage of carrot development

Do not apply within 30 days of harvest

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

When applying to emerged weeds add 2 qt of a nonionic surfactant (NIS) or wetting agent (approved for intended use) to 100 gal of spray mixture (0.5%) v/v or 1 gal of a non-phytotoxic crop oil concentrate (COC) containing 15-20% approved emulsifier to 100 gal of spray mixture (1% v/v).

**Rotational Crops**

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on carrots: cabbage, carrots, celeriac, celery, Chinese celery, Florence fennel, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

---

**CILANTRO**

---

Caparol 4L can be applied post planting, preemergence to cilantro. Read and follow all directions for use in cilantro.

**Use Precautions for Preemergence Applications in Cilantro**

Make one application of Caparol 4L at the rate of 2 – 3.2 pt/A.

Use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and soils high in organic matter.

Do not exceed 3.2 pt/A of Caparol 4L per crop cycle.

Do not apply within 30 days of harvest.

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

Do not use on sand or loamy sand soil.

**Rotational Crops**

The following crops may be seeded 5 months after applying no more than 3.4 pt Caparol 4L per acre on cilantro: cabbage, carrots, celeriac, celery, Chinese celery, Florence fennel, cilantro, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Caparol® 4L is a trademark of Syngenta

Section 24(c) Registrant:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419-8300

EPA SLN No: NY-140007

Label Code: NY0620071AA1014