

RESTRICTED USE PESTICIDE**Toxic to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE
ONLY IN
CALIFORNIA**

THIS PRODUCT BULLETIN CONTAINS NEW OR
ADDITIONAL DIRECTIONS FOR USE WHICH ARE
RECOMMENDED BY FMC CORPORATION UNDER
FIFRA SECTION 2(ee)

**IT IS A VIOLATION OF FEDERAL LAW TO
USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH IT LABELING**

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Strawberries (0 Day PHI)	<i>Vinegar Flies (adults)</i> (<i>Drosophila spp.</i> Including Spotted-wing <i>Drosophila</i>)	8 - 32 oz / acre.

DIRECTIONS FOR USE: Through coverage is essential to achieve control. Use a minimum spray volume of 5 gallons by air or 50 gallons by ground to obtain full coverage of foliage. Multiple applications may be required.

RESTRICTIONS:

Do not apply more than 80 ounces product (0.5 lbs active ingredient) per acre per season. Harvesters and other personnel performing tasks with all-day foliage contact in treated fields within five (5) days of application must wear long-sleeved shirt, long pants, and shoes plus socks.

Following treatment of strawberry fields at rates of Brigade® WSB greater than 0.1 lb ai/acre, harvesters must wear gloves for five (5) days following application.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED
BRIGADE® WSB INSECTICIDE/MITICIDE LABEL (EPA REG. NO. 279-3108) MUST BE FOLLOWED.
THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF
PESTICIDE APPLICATION.**



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Mustang and FMC - FMC Trademarks
©2009 FMC Corporation