RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN OREGON

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

For Agricultural or Commercial Use Only

CROP	PEST	RATE OF APPLICATION
Bushberries including: blueberry; Currant; Elderberry; Gooseberry; Huckleberry Caneberries Strawberries Grape Pear	Vinegar Flies (adults) (Drosophila spp.)	16 oz / acre. (0.1 lb ai/A)

DIRECTIONS FOR USE: Thorough coverage is essential to achieve control. Apply by ground or air using sufficient spray volume to obtain full coverage of foliage. Multiple applications may be required, refer to container label for limits on amount allowed for each crop per season.

BUSHBERRIES (1 Day PHI): Use a minimum spray volume of 20 gallons by ground and 2 gallons by air.

CANEBERRIES (3 Day PHI): Use a minimum spray volume of 50 gallons by ground and 10 gallons by air.

STRAWBERRIES (0 Day PHI): For ground application, use a minimum spray volume of 50 gallons per acre for concentrate spray; for aerial application, use a minimum of 5 gallons per acre.

GRAPES (30 Day PHI): Use a minimum spray volume of 25 gallons by ground and 10 gallons by air.

PEARS (14 Day PHI): Use a minimum spray volume of 50 gallons by ground and 10 gallons by air.

RESTRICTIONS:

Refer to Brigade WSB section 3 label for restrictions.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED BRIGADE WSB INSECTICIDE LABEL (EPA REG. NO.279-3108) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

