



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN THE STATE OF CALIFORNIA

ASSAIL® 30SG INSECTICIDE EPA Reg. No. 8033-36-70506

Additional Use Directions for Fruiting Vegetables

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

PEST	DOSAGE PER ACRE		USE DIRECTIONS
	Pounds Active	Ounces Assail 30SG Insecticide	
Leafhopper	0.047-0.075	2.5-4.0	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

RESTRICTIONS:

- For any of the pests listed above, use the high rate within the listed rate range under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 7 days before harvest (PHI = 7 days).
- Do not exceed a total of 0.3 lbs. acetamiprid active ingredient (16.0 ozs. of ASSAIL 30 SG product) per acre per growing season including any pre-transplant applications of acetamiprid (maximum pre-transplant application rate of acetamiprid is 0.15 lb. ai/A).
- There are no rotation crop plantback restrictions for this product.

Rev. 05/12/2014