

FIFRA SECTION 24(C) SPECIAL LOCAL NEED SUPPLEMENTAL LABEL FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

This label is valid until December 31, 2019, or until otherwise amended, withdrawn, or canceled.

GROUP	21	INSECTICIDE
-------	----	-------------



ACTIVE INGREDIENT: PYRIDABEN	% By Wt.
[2-tert-butyl-5-(4-tert-butylbenzylthio)-4-chloropyridazin-3(2H)-one]	75.0%
OTHER INGREDIENTS:	25.0%
	Total 100.0%

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No 81880-4-10163 / EPA SLN NO. OR-120003 For Control of Pear Psylla Nymphs

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

POLLINATOR PROTECTION: This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Apply Nexter Miticide/Insecticide between late evening and early morning only.

CROP	RATE	PEST	COMMENTS
Pears (including oriental)	10.67 – 16 ounces / acre*	Pear Psylla	Apply Nexter using conventional dilute or concentrate ground sprayers calibrated to deliver enough water to ensure thorough coverage of foliage. Spray volume will vary with the size and number of trees per acre and foliage density. Nexter must be applied to each row for maximum coverage. Pear psylla numbers can increase rapidly; begin application before damaging numbers are reached. Applications may be made early from cluster bud through petal fall to control eggs, early pear psylla instars. Nexter is also effective when applied after petal fall as psylla populations begin to build.

* Water Soluble Bag Acre Conversions (Acres per WSB) for Various Application Rates of Nexter

Nexter Application Rate (rate per acre) oz	Acres Treated per Water Soluble Bag (3.3oz / WSB)
10.67	0.31
11	0.30
12	0.28
13	0.25
14	0.24
15	0.22
16	0.21

To calculate the number of water soluble bags for your spray mix:

1. Determine the number of acres to be treated.
2. Locate proper application rate in the crop-specific section.
3. Divide the total number of acres to be treated by the acres treated per water soluble bag corresponding to the desired rate. (See table above for broadcast application rates and corresponding acres per bag.)
4. ***Restriction:** Do not apply more than 19 Water Soluble Bags (62.7 oz) per 4 acres.

Sample Calculations:

- If the desired application rate is 16 oz/acre, then acres treated/bag = 0.21
- Assuming 17 acres to be treated, 17 acres divided by 0.21 acres/bag = 80 bags.

Note: If the resulting number of bags is not a whole bag, round down to the nearest whole number of bags and check to make sure that the resulting number of acres/bags falls within the desired rate range for the application.

RESTRICTIONS / PRECAUTIONS:

- Do not apply within 28 days of harvest (PHI).
- Do not make more than 1 application per season.
- Do not apply by air.
- For use under this SLN label do not apply this product through any type of irrigation system.
- This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Application early in the morning or at dusk is suggested.

Drift: Do not apply Nexter when weather conditions favor drift to surface water. Do not apply within 110 feet upwind of surface water or when windspeed is above 8 mph. Do not apply during a temperature inversion.

24(c) Registrant: Canyon Group
C/O Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Distributor: Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569