RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG. NO. 279-3426

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN Georgia, Illinois, Delaware, Maine, Maryland, Massachusetts, Michigan, New Jersey, North Carolina, Ohio, South Carolina, Virginia, West Virginia and Wisconsin

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

CROP	PEST	RATE OF APPLICATION
Berries Crop Group including: blackberry; loganberry; red and black raspberry; blueberry, highbush and lowbush; currant; elderberry; gooseberry; huckleberry; and cultivars and/or hybrids of these.	Controls The Following Pests: Cherry Fruitworm, Cranberry Fruitworm, Cranberry Weevil, Japanese Beetle, Plum Curculio, and Scale (crawlers).	4.0 fl oz / acre
	Suppression Only: Green Aphid, Blueberry Aphid, Blueberrry Maggot	

DIRECTIONS FOR USE: Through coverage is essential to achieve control. Apply using sufficient spray volume to obtain full coverage, use a minimum spray volume of 20 gallons by ground or 2 gallons by air. Multiple applications may be required

Pre-harvest interval is 1 day.

RESTRICTIONS:

Do not apply more than 24 fluid ounces product (0.15 lbs active ingredient) per acre per season.

Do not make applications less than 7 days apart.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG MAX INSECTICIDE LABEL (EPA REG. NO. 279-3426) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.