SUPPLEMENTAL LABEL

FOR USE ON ONION, BULB AND GREEN (SUBGROUPS 3-07A and 3-07B)



KEEP OUT OF REACH OF CHILDREN CAUTION

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

DIRECTIONS FOR USE ON ONION, BULB AND GREEN (subgroups 3-07A and 3-07B)

CROP	PEST	RATE	COMMENTS	
Bulb onion,	Flea Beetles	FOLIAR:	Higher water volumes provide improved insect control.	
includes:	Grasshoppers	3.5 to 7 fl oz/A		
Daylily, bulb	Leafhoppers	(0.090 to 0.180 lb ai/A)	Begin applications when first pest activity is noticed or when	
Fritillaria, bulb	Stink bugs	FOLIAR:	insects reach threshold levels per State and County Extension	
Garlic, bulb	Leafminers	5.25 to 7 fl oz/A	Service recommendations. Repeat as needed to maintain control,	
Garlic, Great-	Thrips	(0.135 to 0.180 lb ai/A)	but not more often than every 7 days. For best results, time	
headed, bulb	Whiteflies		application before a damaging population becomes established.	
Garlic, serpent, bulb	Leafminers	SOIL:	application before a damaging population becomes established.	
Lily, bulb Onion, bulb	Thrips	8.75 to 10.5 fl oz/A	II. dan account management the bighter management design	
Onion, Chinese,	Whiteflies	(0.225 to 0.270 lb ai/A)	Under severe pest pressure, use the higher specified rates.	
bulb	· · · · · · · · · · · · · · · · · · ·			
Onion, pearl			Restriction: Do not apply to vegetables grown for seed.	
Onion, potato, bulb				
Shallot, bulb			The rate applied affects the length of control. Use the high rate	
Cultivars, varieties			where infestations occur later in crop development, or where pest	
and/or			pressure is continuous.	
hybrids of these				
			SCORPION can be mixed and/or alternated with commonly used	
Green onion,			insecticides, such as <i>Knack</i> , to improve the length of control	
includes:			and/or achieve better knockdown of pests	
Chive, fresh leaves	Note: Regardless of ap	plication method of SCORPI	ON do not exceed 15 fl oz/A (0.383 lb ai/A) per crop season.	
Chive, Chinese, fresh leaves				
Elegans hosta	Foliar Application			
Fritillaria leaves	Apply with air or ground equipment in adequate water for uniform coverage (A minimum of 5 gals/A by air or 20 gals/A by			
Kurrat	ground).			
Leady's leek	• Do not apply SCORPION within one (1) day of harvest.			
Leek	• Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per season.			
Leek, wild	Sail Application			
Onion, Beltsville	Soil Application • See conversion chart for linear application plant application rates.			
bunching	Apply with ground equipment in adequate water for uniform coverage (A minimum of 10 gals/A).			
Onion, fresh	Apply SCORPION at planting or immediately after transplanting.			
Onion, green	• Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per season.			
Onion, macrostem	Do not apply more than	a total of 10.0 ii 027 (0) 0001	(1 1011 (0.270 10 df) por doro por obdeorii.	
Onion, tree, tops	Apply specified dosage in	Apply specified dosage in sufficient carrier volume to insure uniform application and incorporate into the soil using one of		
Onion, Welsh, tops	the following methods: 1. In a narrow band centered on the plant row in the bedding operation just prior to planting. For best results band width			
Shallot, fresh leaves				
Cultivars, varieties and/or	should be 2" or less and placed 1 to 2" below the seed depth.			
hybrids of these	2. In-furrow spray at or below seed level or a narrow surface band above the seedline during planting. For surface-banded			
Hybrids of these	applications incorporate to a depth of 1-1/2" with sufficient irrigation within 24 hours to insure satisfactory insect control.			
		rench, transplant drench or hill drench. Applications should be made with sufficient water to insure		
	incorporation into the root zone.			
	4. As a sidedress immediately after transplanting operations are finished. Applications should be placed within 2 to			
	side of each row and incorporated 1 or more inches deep. Applications should be made to each row if there are two rows per bed.			
	per bed.			

EPA Reg. No. 10163-317 EPA Est. No.

