SUPPLEMENTAL LABEL FOR USE ON WATERCRESS



KEEP OUT OF REACH OF CHILDREN CAUTION

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

DIRECTIONS FOR USE ON WATERCRESS

CROP	PEST	RATE	COMMENTS
Watercress	Cucumber Beetle Sharpshooters	FOLIAR: 3.5 to 7 fl oz/A	Higher water volumes provide improved insect control.
	Leafhoppers Fleabeetles	(0.090 to 0.180 lb ai/A)	Begin applications when first pest activity is noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat as needed to maintain control, but not more often than every 7 days. For best results, time application before a damaging population becomes established.
	Aphids (suppression) Stinkbugs Whiteflies	Stinkbugs 5.25 to 7 fl oz/A Whiteflies (0.135 to 0.180 lb ai/A)	
	Thrips		Under severe pest pressure, use the higher specified rates.
			The rate applied affects the length of control. Use the high rate where infestations occur later in crop development, or where pest pressure is continuous.
			SCORPION can be mixed and/or alternated with commonly used insecticides, such as <i>Danitol</i> or <i>Knack</i> , for better knockdown and/or improved control of pests.

Foliar Application

- Apply with air or ground equipment in adequate water for uniform coverage (5 to 10 gals/A by air or 50 to 300 gals/A by ground).
- Do not apply SCORPION within one (1) day of harvest.
- Interval between application cannot be less than 7 days
- Do not apply more than a total of 14 fl oz/A of SCORPION (0.360 lb. ai) per acre per crop season.

EPA Reg. No. 10163-317 EPA Est. No.



Distributed By: Gowan Company PO BOX 5569 Yuma, AZ 85366