



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

## ADMIRE<sup>®</sup> PRO Systemic Protectant

Provide for alternative application methods for pest management in HI Banana and Tropical Fruit

EPA Reg. No. 264-827

# PRODUCT BULLETIN

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF HAWAII

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

### BANANA and PLANTAIN

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers	7.0 – 14.0
Pests Suppressed	
Scales	7.0 – 14.0

### NOTES AND RESTRICTIONS

Pre-Harvest Interval (PHI): **0 days**

Maximum ADMIRE PRO allowed per season: **14.0 fluid ounces/Acre** (0.5 lb AI/Acre).

### Applications

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation;
3. Basal, soil drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

## TROPICAL FRUIT

Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids Avocado lace bug Leafhoppers Whiteflies	10.5 – 14.0
Pests Suppressed	
Scales Thrips (foliage feeding thrips only)	14.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>6 days</b> Maximum ADMIRE PRO allowed per season: <b>14.0 fluid ounces/Acre</b> (0.5 lb AI/Acre). Do not apply pre-bloom or during bloom or when bees are actively foraging <b>Applications</b> Apply specified dosage in one of the following methods: 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation; 3. Basal, soil drench in sufficient water to insure incorporation into the root-zone followed by irrigation.	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*