

Supplemental Label



For use in safflower

This supplemental label expires March 1, 2018, and must not be used or distributed after this date.

Active Ingredient*:

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine 38.7%

Other Ingredients: 61.3%

Total: 100.0%

*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

EPA Reg. No. 241-418

CAUTION/PRECAUCIÓN

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl® H₂O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H₂O** before applying.
- Use of **Prowl H₂O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H₂O**.

Product Information

Prowl H₂O is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H₂O** container label for a complete list of weeds controlled.

Application Instructions

Prowl H₂O may be applied to safflower by ground or air.

Plant safflower 1-1/2 inches to 2-inches deep and completely cover with soil.

Use Methods and Timings

Preplant Incorporated. In all states, apply within 60 days of planting and incorporate.

Preplant Incorporated

(Fall Applications Only in Minnesota, North Dakota, and South Dakota). Apply **Prowl H₂O** and immediately incorporate in late fall before planting safflower the following spring. Apply **Prowl H₂O** in the late fall when soil temperatures are 45° F or below but before the ground freezes. **DO NOT** apply when air temperature is below 45° F.

Before safflower planting in the spring, fields treated with **Prowl H₂O** should receive at least one shallow additional incorporation. Spring incorporation should be at an angle to the last tillage operation.

Preemergence. Apply **Prowl® H₂O** herbicide at planting or up to 2 days after planting.

Preemergence application of **Prowl H₂O** to safflower may increase the likelihood of crop injury, especially when crops are grown in stress situations, such as compacted soils. Decreased herbicide performance compared to preplant incorporated applications may also result from a preemergence application. If dry conditions with limited precipitation exist or unseasonably cool temperatures following planting are forecast, apply **Prowl H₂O** before planting and mechanically incorporate with tillage. **Prowl H₂O** may be applied preemergence in conventional tillage safflower, **except in the state of California.**

No-till Safflower. **Prowl H₂O** may be applied at 3.0 pts/A up to 30 days before planting (preplant) to immediately after planting (preemergence). **DO NOT** use in California.

Use Rates

Preplant Incorporated or Preemergence (Conventional Tillage)

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹	
		< 3% Organic Matter > 3% (pts/A)	
Coarse	1.5	2.0	2.0
Medium	2.0	2.5	3.0
Fine	3.0	3.0	3.0
¹ See Restrictions and Limitations on product container label for map of specific states.			

Preplant Incorporated (Fall) Application¹

Soil Texture	< 3% Organic Matter > 3% (pts/A)	
Coarse	2.5	2.5
Medium	3.0	3.5
Fine	3.5	3.5
¹ For use in Minnesota, North Dakota, and South Dakota only		

Restrictions and Limitations (All Tillage Types)

- **DO NOT** apply **Prowl H₂O** postemergence.
- **DO NOT** exceed the highest rate per acre for any soil type.
- **DO NOT** feed forage or graze livestock in treated safflower fields.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Prowl is a registered trademark of BASF.

© 2015 BASF Corporation
All rights reserved.

000241-00418.20150223.NVA 2015-04-195-0029
Based on: NVA 2013-04-195-0111

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry