

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **CYONARA LP** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **CYONARA LP** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

CYONARA LP

Lawn & Perimeter Insect Control Granules

EPA Reg. No. 53883-195

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

In case of an emergency with this product contact SafetyCall® International at (866) 897-8050.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Keep children and pets off treated area until product has been watered in and lawn is dry.

Restrictions:

- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the labels of these products.
- Do not apply more than 0.36 lbs. of the a.i. (800 pounds of CYONARA LP) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

LAWN AND PERIMETER PEST CONTROL

Cyonara LP insecticide may be used for applications to maintain outdoor areas where turf and ornamental ground covers are grown such as residential landscaped areas and non residential landscapes around institutional, public, commercial and institutional buildings, parks, recreational areas, and athletic fields. Broadcast Cyonara LP Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. The application rates listed in the **Lawn and Perimeter Use Rate** table will provide excellent control of the respective pests under typical conditions.

NOTE:

- Use higher rate for longer length of control.
- Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide.
- Make no more than 6 general applications per year. More frequent treatments should be limited to spot treatments.
- Uniform coverage is necessary to provide the most effective level of control.
- Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.
- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Recommended Application Rates for Lawn and Perimeter Use

	PEST	A.I. lb per acre	Cyonara LP Rate Per 1000 ft ²	Cyonara LP Rate Per Acre
Band Treatment Around Foundation	Ants Crickets Firebrats Mites Sowbugs Beetles Earwigs Millipedes Silverfish	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Ants Armyworms Centipedes Crickets Cutworms Earwigs Grasshoppers Green June Beetles Japanese Beetle (Adult) Leafhoppers Millipedes Mites Pillbugs Sod webworms Spider Mites (Twospotted, Spruce) Spittlebugs Springtails and Ticks (including Ticks which may transmit Lyme Disease)	0.04-0.06	2 - 3 lbs.	88 – 131 lbs.

Overall Lawn Application	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	0.06-0.078	3 - 4 lbs	131 - 176 lbs.
Spot Treatment	Ants		Apply 1-2 teaspoons around each ant hill	

*Suppression only.

SPECIFIC USE INSTRUCTIONS

Treatment Around Foundations

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Nuisance), Fleas, Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs

Application Rate: Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 5' x 200') oz. or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').

Application Instructions: For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.

Overall Lawn Application

Chinch Bug and Mole Cricket Control -

For **chinch bug control**, apply at higher rates during the summer when both nymphs and adults may be present.

For **mole crickets** apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.

Spot Treatment

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Nuisance)

Mound Treatment: Apply 1-2 teaspoons over and around each ant hill.

NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle an open container in a manner as to prevent spillage.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-refillable container. Do not re-use or re-fill this container. Offer for recycling if available.

Completely empty container. Dispose of container at an approved waste disposal facility or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed, or drinking water.

Spills: If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away.

Read the “NOTE” on the label for Cyonara LP before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041