

SUPPLEMENTAL LABELING

GROUP 11 FUNGICIDE

EVITO®480 SC Fungicide

For control of certain diseases in corn (field, sweet and hybrid seed corn), fruiting vegetables and leaf petiole vegetables, listed low growing berries, peanuts, potato and other tuberous and corm vegetables, soybean, and wheat.

ACTIVE INGREDIENT:

Fluoxastrobin: [(1*E*)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] 5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime]**40.3%**

INERT INGREDIENTS:.....**59.7%**

TOTAL: 100.0%

This product contains 4 pounds of fluoxastrobin per gallon (480 g per liter)

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label for

EVITO 480 SC Fungicide (EPA Reg. No. 66330-64).

This supplemental label must be in the possession of the user at the time of application.

GENERAL INFORMATION

EVITO™ 480 SC Fungicide is a broad-spectrum fungicide for the control of certain diseases in corn (field, sweet and hybrid seed corn), fruiting vegetables and leaf petiole vegetables, listed low growing berries, peanuts, potato and other tuberous and corm vegetables, soybean, and wheat. EVITO™ 480 SC Fungicide works by interfering with respiration in plant-pathogenic fungi, and is a potent inhibitor of spore germination and mycelial growth.

AERIAL APPLICATION

Corn (Field, Sweet and Hybrid Seed Corn, listed Low Growing Berries, Soybean, Tuberous and Corm Vegetables, and Wheat only)

For aerial application a minimum of 5 gallons/A is recommended. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Aerial applications made to dense canopies may not provide sufficient coverage of lower leaves to provide proper pest control.

CHEMIGATION

Corn (Field, Sweet and Hybrid Seed Corn, Fruiting Vegetables, Leaf Petiole Vegetables, Tuber and Corm Vegetables, Soybean, and Wheat only)

Apply EVITO™ 480 SC Fungicide only through sprinkler type irrigation systems, including center pivot, microjet, wheel lines, lateral move, side roll, or overhead solid set irrigation systems. Do not apply EVITO™ 480 SC Fungicide through any other type of irrigation system.

USE DIRECTIONS FOR SPECIFIC CROPS

EVITO 480 SC Fungicide provides control or suppression of several important diseases of corn (field, sweet and hybrid seed corn), fruiting vegetables and leaf petiole vegetables, listed low growing berries, peanuts, potato and other tuberous and corm vegetables, soybean, and wheat. When reference is made to disease suppression, suppression can mean either erratic control from good to fair, or consistent control at a level below that obtained with the best commercial disease control products.

SOILBORNE/SEEDLING DISEASE CONTROL

(Only for Corn (field, sweet and hybrid seed corn), Soybean, and Low-growing berry (crop subgroup 13-07G))

EVITO 480 SC Fungicide can provide control of many soilborne diseases if applied early in the growing season. Specific applications for soilborne diseases include in-furrow applications or banded applications applied over the row, either shortly after plant emergence or during herbicide applications or cultivation. These applications will provide control of pre- or post-emergence damping off and diseases that infect plants at the soil-plant interface. The use of either type of application depends on the cultural practices in the region. In some locations, one type of application may provide better disease control than the other, depending on the timing of the disease epidemic. Seedling diseases are generally controlled by in-furrow applications while banded applications are more effective against soilborne diseases that develop later in the season. Consult your local expert to get some guidance regarding application type.

CORN (Field, Sweet and Hybrid Seed)

Disease Control	Rate to Use	Application Timing and Resistance Management
Rust, common <i>(Puccinia sorghi)</i> Rust, southern <i>(Puccinia polyspora)</i> Anthracnose leaf blight <i>(Colletotrichum graminicola)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i> Northern corn leaf blight <i>(Setosphaeria turcica)</i> Northern corn leaf spot <i>(Cochliobolus carbonum)</i> Southern corn leaf blight <i>(Cochliobolus heterostrophus)</i> Eye Spot <i>(Aureobasidium zeae)</i>	For Field and Hybrid Seed Corn Apply: 2.0 to 5.7 fl oz/A (0.063 to 0.18 lbs ai/A) For Sweet Corn Apply: 2.0 to 3.8 fl oz/A (0.063 to 0.12 lb ai/A)	Apply a maximum of two applications preventatively, (with the final application no later than the R4 early dough stage). For optimum results, begin applications preventively and continue as needed on a 7 to 10-day interval on field and seed corn and a minimum 14-day interval on sweet corn. Use the higher rates and shorter interval when disease pressure is high. Resistance Management: Do not make more than two (2) sequential applications of Evito before alternating to a labeled fungicide with a different mode of action for at least one (1) application.
SOILBORNE DISEASES CONTROL		
Rhizoctonia Root and stalk rot <i>(Rhizoctonia solani)</i>	0.16-0.24 fl oz/1,000 row feet	For soilborne/seedling disease control, see directions and rates under GENERAL INFORMATION section.

RESTRICTIONS AND OTHER INFORMATION:

Field and Hybrid Seed Corn

- Do not apply more than 11.4 fl oz (0.36 lb ai)/A of EVITO™ 480 SC Fungicide per acre per year (including from an in-furrow or banded application).
- There is a maximum number of 2 applications per season, and a minimum interval of 7 days between applications.
- EVITO 480 SC Fungicide may also be applied through chemigation or by air.
- Do not apply EVITO™ 480 SC Fungicide after the R4 stage (early dough).
- Do not apply EVITO™ 480 SC Fungicide within 30 days of harvest.
- Do not use the 0.24 fl oz/1,000 row feet rate on rows spaced narrower than 22 inches.
- Do not use the 0.16 fl oz/1,000 row feet rate on rows spaced less than 15 inches.
- For twin rows spaced 7.5 to 8" apart on 30 inch centers, use the 0.16 fl oz/1,000 row feet rate for 15" rows only.

Sweet Corn

- Do not apply more than 15.2 fl oz (0.48 lb ai)/A of EVITO™ 480 SC Fungicide per acre per year (including from an in-furrow or banded application).
- There is a maximum number of 4 applications per season, and a minimum interval of 14 days between applications.
- EVITO 480 SC Fungicide may also be applied through chemigation or by air.
- EVITO 480 SC Fungicide may be used with an adjuvant to increase coverage.
- Do not apply EVITO 480 SC Fungicide within 7 days of harvest.

- Do not apply EVITO 480 SC Fungicide within 23 days of use of stover for feed.

WHEAT

Disease Control	Rate to Use	Application Timing and Resistance Management
Leaf rust <i>(Puccinia recondita f. sp. tritici)</i> Stripe rust <i>(Puccinia striiformis)</i> Stem rust <i>(Puccinia graminis)</i> Septoria leaf and glume blotch <i>(Septoria tritici, Septoria nodorum)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	2.0 to 4 fl oz/A (0.063 to 0.12 lb ai/A)	For optimum results, begin applications preventively and continue as needed on a 14 to 21 day interval. Use the higher rates and shorter interval when disease pressure is high Resistance Management: Do not make more than two (2) sequential applications of EVITO™ 480 SC Fungicide before alternating to a labeled fungicide with a different mode of action for at least one (1) application. Apply prior to disease development from Feekes 5 (Zadok's 31) up to late head emergence at Feekes 10.5 (Zadok's 59).
Powdery Mildew <i>(Erysiphe graminis)</i>	2.5 to 4 fl oz/A (0.079 to 0.12 lb ai/A)	

RESTRICTIONS AND OTHER INFORMATION:

- Do not apply more than 8 fl oz (0.24 lb ai) of EVITO™ 480 SC Fungicide per acre per year.
- There is a maximum number of 2 applications per season, and a minimum interval of 14 days between applications.
- EVITO 480 SC may also be applied through chemigation or by air.
- Do not apply EVITO™ 480 SC Fungicide within 40 days of harvest for grain and straw.
- Do not apply EVITO™ 480 SC Fungicide within 7 days of harvest for forage and hay.
- Do not apply later than Feekes growth stage 10.5.
- Make no more than one application prior to harvest of wheat forage.



Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

EVITO is a registered trademark of Arysta LifeScience North America, LLC
Arysta LifeScience and the Arysta LifeScience logo registered trademarks of Arysta LifeScience Corporation