



GROUP 11 FUNGICIDE

EVITO®480 SC Fungicide

Supplemental Label to Add Directions for Use on Squash/Cucumber Subgroup 9B.

For Agricultural Uses

ACTIVE INGREDIENT:

Fluoxastrobin: [(1*E*)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime] **40.3%**

OTHER INGREDIENTS: **59.7%**

This product contains 3.69 pounds of fluoxastrobin per gallon (443 g per liter) **TOTAL: 100.0%**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Before using EVITO 480 SC as permitted according to this supplemental label, read and follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label for

EVITO 480 SC Fungicide (EPA Reg. No. 66330-64).

This supplemental label must be in the possession of the user at the time of application.

PRODUCT INFORMATION

EVITO® 480 SC Fungicide is a broad-spectrum fungicide for the control of certain diseases in squash /cucumber subgroup 9B. EVITO® 480 SC Fungicide works by interfering with respiration in plant-pathogenic fungi, and is a potent inhibitor of spore germination and mycelial growth.

SQUASH/CUCUMBER SUBGROUP 9B*

Including: Chayote, Chinese waxgourd (Chinese preserving melon), Cucumber, Gherkin, Gourd, edible (includes hyotan, cucuzza, h echima, Chinese okra, balsam apple, balsam pear, bitter melon, Chinese cucumber), Pumpkin, Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini), and Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash).

Foliar Disease Control	Rate to Use	Application Timing and Resistance Management
<i>Alternaria</i> blight <i>(Alternaria cucumerina)</i> Anthrachnose <i>(Colletotricum orbiculare)</i> Belly rot <i>(Rhizoctonia solani)</i> <i>Cercospora</i> leaf spot <i>(Cercospora citrulina)</i> Downy mildew <i>(Pseudoperonospora cubensis)</i> Gummy stem blight <i>(Didymella bryoniae)</i> <i>Microdomium</i> blight <i>(Plectosporium tabacinum)</i> <i>Myrothecium</i> canker <i>(Myrothecium roridum)</i> <i>Plectosporium</i> blight <i>(Plectosporium tabacinum)</i> Powdery mildew <i>(Sphaerotheca fuliginea,</i> <i>Erysiphe cichoracearum)</i> Target leaf spot <i>(Corynesporium cassicola)</i>	3.0 to 5.7 fl oz/A (0.09 to 0.18 lb ai/A)	<p>Evito 480SC Fungicide should be used in conjunction with good crop management practices and integrated into an overall disease management strategy.</p> <p>For optimum results, begin applications preventively and continue as needed on a 7 to 14-day interval. Higher application rates should be used when disease is severe. For belly-rot control, make the first application at the 1-3 leaf crop stage, followed by a second application 10-14 days later, or at vine tip-over, whichever occurs first.</p> <p>Resistance Management: Alternate every application of Evito 480SC Fungicide with at least one application of another registered, non-Group 11 fungicide.</p> <ul style="list-style-type: none"> Do not make more than 4 applications of Evito 480 SC Fungicide and or a Group 11 fungicide per season. Do not use Evito 480 SC Fungicide for control of gummy stem blight where resistance to Group 11 (QoI) fungicides exists. Additional information on resistance management can be found in the General Information section of this label.
Soilborne Disease Control	Rate to Use	Application Timing and Resistance Management
Root rot <i>(Rhizoctonia solani)</i>	0.16-0.24 fl oz/1,000 row feet	See general application information

*Not approved for use on squash/cucumber subgroup 9B in California.

RESTRICTIONS AND OTHER INFORMATION:

- Do not apply more than 22.8 fl oz (0.66 lb ai) of EVITO 480 SC Fungicide per acre per year.
- There is a maximum number of 4 applications per season, and a minimum interval of 7 days between applications.
- EVITO 480 SC Fungicide may also be applied through chemigation on subgroup 9B crops.
- Do not apply to subgroup 9B crops grown in a greenhouse.
- Do not apply EVITO 480 SC Fungicide within 1 day of harvest.
- Do not tank mix Evito 480 SC Fungicide with EC-based insecticides, or the following products, as this may increase the risk of crop injury under certain environmental conditions: malathion, Kelthane®, Thiodan®, Phaser®, Lannate®, Lorsban®, M-Pede® or Botran®, as crop injury may occur.
- EVITO 480 SC Fungicide may be used with a Non-Ionic Surfactant (NIS).
- Refer to the APPLICATION GUIDELINES for information on tank mixes and the use of adjuvants.
- Do not use the 0.24 fl oz/1,000 row feet rate on rows spaced narrower than 22 inches.
- Do not use the 0.16 fl oz/1,000 row feet rate on rows spaced less than 15 inches.
- For twin rows spaced 7.5 to 8" apart on 30 inch centers, use the 0.16 fl oz/1,000 row feet rate for 15" rows only.



Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

EVITO is a registered trademark of Arysta LifeScience North America, LLC
Arysta LifeScience and the Arysta LifeScience logo registered trademarks of Arysta LifeScience Corporation
SuppAD081811_083111