

TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF VENERATE™ XC Bioinsecticide ON CHEMIGATION TREATMENT FOR POTATOES

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

**VENERATE™ XC**

**EPA Reg. No. 84059-14**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Potatoes	<b>Control</b> Armyworms Artichoke plume moth European corn borer Loopers Psyllids  <b>Suppression</b> Aphids Potato aphid Stink bugs Whiteflies	Chemigation	2–4 quarts per acre	<p><b>CHEMIGATION USE</b>  Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand moved irrigation systems. Do not apply this product through any other type of irrigation systems. Do not connect an irrigation system (including greenhouse systems) used for pesticide applications to a public water system. Do not use reclaimed water for application of this product.</p> <p><b>Spray preparation</b>  First prepare a suspension of VENERATE™ XC in a mix tank. Fill tank ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of VENERATE™ XC, and then the remaining volume of water. Then set the sprinkler to deliver a minimum of 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of VENERATE™ XC into the irrigation water line so as to deliver the desired rate per acre. Inject the suspension of VENERATE™ XC with a positive displacement pump into the main line after the filter, and ahead of a right angle turn to insure adequate mixing. Any questions on calibration should be directed to your State Extension Service Specialists, to equipment manufacturers or other experts.</p> <p>Do not combine VENERATE™ XC with pesticides, surfactants or fertilizers for application through chemigation equipment unless prior experience has shown the combination physically compatible, effective and non-injurious under conditions of use. VENERATE™ XC has not been fully evaluated for compatibility with all adjuvants or surfactants. It is advisable to conduct a spray compatibility test if a mixture with adjuvants or surfactants is planned.</p>
VENERATE™ XC has a <b>pre-harvest interval (PHI)</b> of <b>0 days</b> . Do not enter or allow worker entry into treated areas during the <b>restricted-entry interval (REI)</b> of <b>4 hours</b> .				

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

VENERATE™ is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2014

Marrone Bio Innovations, Inc.

2121 Second Street, Suite B-107

Davis, California 95618

Telephone (530) 750-2800

[www.marronebioinnovations.com](http://www.marronebioinnovations.com)

**2121 Second Street, Suite B-107 • Davis, CA 95618 • Phone: 530-750-2800**