



Cornerstone[®] 5 Plus Herbicide
EPA Reg. No. 1381-241

PRODUCT BULLETIN

**For Postemergence Applications to Spring Canola
with the Roundup Ready[®] Gene**

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Postemergence Applications to Spring Canola with the Roundup Ready Gene: This product may be applied postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Single application: Apply 12 to 18 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and/or growth reduction. Similar injury may result with applications of more than 12 fl. oz. per acre are applied after the 4-leaf stage.

Sequential applications: Apply 12 fluid ounces per acre to 1 – 3 leaf canola followed by a sequential application at a minimum interval of 10 days, but not later than the 6-leaf stage. Sequential application are recommended for early emerging annual weed and perennial weeds such as Canada thistle and quackgrass or when controlling weeds with multiple application times.

Precautions, Restrictions: No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop applications should not exceed 24 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

Important: Refer to the Cornerstone 5 Plus label for additional information, including weeds controlled and general use precautions and restrictions.

Cornerstone is a registered trademark of Winfield Solutions, LLC
Roundup Ready is a registered trademark of Monsanto Technology, LLC

Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164

03/08/2010