

RESTRICTED USE PESTICIDE DUE TO HIGH ACUTE TOXICITY TO HUMANS

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application; mixing; loading or repair or cleaning of application equipment.

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ LANNATE® SP
INSECTICIDE
FOR USE ON CHINESE CABBAGE
(NAPA, BOK CHOY)**

DUPONT™ LANNATE® SP INSECTICIDE

EPA Reg. No. 352-342

This supplemental label expires on June 1, 2018 and must not be used or distributed after this date.

FOR USE ON CHINESE CABBAGE (NAPA, BOK CHOY)**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ LANNATE® SP must be used in accordance with the directions for use on this label. Always read the entire label affixed to the container (including the Limitation of Warranty and Liability) before applying. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Use of Lannate SP Insecticide, according to this labeling, is subject to the use precautions and limitations, imposed by the label affixed to the container, for LANNATE® SP Insecticide."

IMPORTANT

Before using LANNATE® SP, read and follow all applicable directions; restrictions; and precautions on the label affixed to the container. This supplemental label contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

APPLICATION

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Follow-up treatments of DuPont™ LANNATE® SP should be applied, as needed, to keep pest populations within threshold limits. On most crops, LANNATE® SP should be applied at 5 to 7 day intervals to maintain control.

Use sufficient water to obtain thorough, uniform coverage. Since LANNATE® SP is a fast acting contact insecticide, best results follow direct spraying of the target insect.

Apply the low rates on small plants, small insects and light infestations of insects. Use intermediate rates on large insects and heavier infestations of insects. Use 1 to 3 applications of the highest recommended rate for controlling severe infestations. Thereafter, use the lowest rate possible to maintain control.

© 2015 E. I. du Pont de Nemours and Company, Crop Protection, 1007 N. Market St., Wilmington, Delaware 19898

Crops	Insects	Rate DuPont™ LANNATE® SP Lbs. Per Acre	Last Application -Days To Harvest	REI
Chinese Cabbage (Napa, Bok Choy)	Loopers Beet Armyworm	1/2 - 1*	10	48 hrs
	Do not apply more than 8 pounds of LANNATE® SP/acre/crop. Do not make more than 10 applications/crop. * Minimum of 25 gallons water per acre by ground or 5 gallons by air.			

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.