



Section 2(ee) Recommendation

Date: July 30, 2015
Product: Ridomil Gold® SL
EPA Reg. No. 100-1202
Use: Addition of Late Blight (*Phytophthora infestans*) as a Pest Controlled on Potato
State: Idaho

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

- The user must refer to the federally approved labels and approved Supplemental Label, SCP 1202A-S5 0614, for the above noted product and read and follow all directions for use, restrictions, and precautions.

Crop	Pest	Rate	Instructions
Potato	Late Blight (<i>Phytophthora infestans</i>)	3.2 oz/A (0.1 lb ai)	A tank mix with a labeled rate of mancozeb or chlorothalonil products is required. Begin preventative applications when conditions are favorable for disease. Apply on a 14 day schedule and alternate with an effective non-Group 4 fungicide. Use in conjunction with other management practices such as crop rotation and resistant varieties.

- Do not exceed the equivalent of 0.34 lb ai/A per season of soil-applied and 0.4 lb ai/A per season of foliar applied mefenoxam-containing products.
- Do not apply within 14 days of harvest (14-day PHI).

Ridomil Gold® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2015 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: July 30, 2020

Label Code: ID1202127AC0715